

Global Animal Partnership's 5-Step® Animal Welfare Standards for Chickens Raised for Meat v3.2



About Global Animal Partnership's (G.A.P.) Animal Welfare Certified™ Program

G.A.P.'s Animal Welfare Certified™ program assesses the level of welfare of animals on-farm, during transport and at slaughter. Standards for additional components of production, such as hatchery and breeding, will be developed in the future.

Through its very design as a multi-tiered program, G.A.P.'s Animal Welfare Certified™ Program promotes continuous improvement in animal agriculture; engages a broad spectrum of producers; allows for wider product selection for greater consumer options; provides more detailed information about the level of welfare of farm animals; and better informs consumers about the production systems they may choose to support.

Each set of tiered standards—from Step 1 to Step 5+—has its own requirements that must be met before certification to that particular Step level can be assigned. As the standard-setter, Global Animal Partnership does not conduct audits nor make Step-level certification decisions. Authorized, third-party certification companies perform the audits and issue Step certificates, as appropriate. As such, producers, consumers, and retailers alike can be confident that Step-levels are fair, accurate, and free of conflict of interest.

About G.A.P.'s 5-Step® Animal Welfare Standards for Chickens Raised for Meat

Version 3.0 of the G.A.P.'s 5-Step® Animal Welfare Standards for Chickens Raised for Meat was issued on 3 July 2017, was amended (v3.1) on 3 April 2018, and was updated (v3.2) on 1 May 2020. This version (3.1) replaces all previous versions. Any operations audited on or after 3 April 2018, including those undergoing re-certification, will be audited to this version. The development process included consultation and guidance from animal welfare scientific experts, chicken producers and industry representatives, public comments, and review and approval by the Global Animal Partnership Board of Directors.

After four (4) certification cycles, the standards will be reviewed in full and revised through a process that involves expert guidance from scientists and producers, public comment and field testing, before the draft revision is presented to the Board of Directors for final review and approval. This time frame provides for the standard to be implemented across each of the seasons prior to a thorough review and revision process.

At any time throughout the above-mentioned period, G.A.P. may amend or clarify parts of the standard as issues with implementation arise, new technologies become available, or new scientific findings are made.





About Global Animal Partnership



Global Animal Partnership, a nonprofit charitable organization founded in 2008, brings together farmers, scientists, ranchers, retailers, manufacturers, foodservice and animal advocates—a diverse group with the common goal of seeking to improve the welfare of farm animals.

For more information, contact us at info@globalanimalpartnership.org or 877.427.5783.

Program Overview

Chickens raised in Step 1 systems typically live in a stationary housing structure and are provided with space to express natural behavior and enrichment in their environment. In Step 2, they also typically live in an indoor environment but with natural light and are provided with different types of enrichments. Chickens in Step 3 systems have access to the outdoors, when seasonal conditions allow, with provisions that encourage ranging and foraging when outdoors. In Step 4, chickens live on pasture, with access to housing; during winter, chickens may be taken off pasture but must have daily access to the outdoors. In Step 5 and Step 5+, chickens live continuously outdoors on pasture and may only be removed during extreme weather conditions. At Step 5+, chickens are also required to spend their lives from placement on a single farm, and are slaughtered at a mobile or on-farm slaughter facility.

Step Level	Marketing Claims	Description of System	Management Highlights
	No cages, no crates, no crowding	Chickens in Step 1 systems live in a stationary housing structure and are provided space to express natural behavior and given access to added enrichments.	Chickens are typically housed indoors and must meet a maximum stocking density of 6.0 lbs/ft ² by 1 July 2020. Producers are required to manage the environment to maintain litter, air quality and provide chickens with environmental enrichment. Environmental enrichments are materials that are provided to chickens to add complexity to their environment and encourage the expression of natural behavior (such as pecking, scratching, exploration and play behavior).
	Enriched environment	Chickens in Step 2 systems live in an indoor environment with at least two different types of enrichment and natural light (by 2022).	At Step 2, chickens must have two different types of enrichments to further encourage the expression of natural behaviors and the operation must be progressing towards the provision of natural light.
	Enhanced outdoor access	Chickens in Step 3 systems have seasonal outdoor access.	Chickens live in a stationary housing structure with natural light with seasonal outdoor access. Outdoor access is defined as an outdoor area but does not have to be pasture. Outdoor areas must contain features that increase the complexity of the environment. Indoor areas must have two different types of enrichments and natural light.
	Pasture centered	Chickens in Step 4 systems live on pasture; during winter, chickens may be housed with continuous access to the outdoors.	This step requires breeds of chickens with higher welfare outcomes that live on pasture from the age of 4 weeks with access to housing or shelter. During winter, chickens may be taken off pasture but must have daily access to the outdoors. Pasture includes access to rangelands, grassland, planted pastures, managed pastures, wooded areas, and harvested crop areas.

	Animal centered; no physical alterations	Chickens in Step 5 systems live continuously on pasture and may only be housed during extreme weather conditions.	This Step requires breeds with higher welfare outcomes that have the ability to perch and live continuously on pasture from the age of 4 weeks. This step also requires confirmation that the breeding stock meet specific welfare criteria.
	Animal centered; entire life on same farm	Chickens in Step 5+ systems live continuously on pasture and may only be housed during extreme weather conditions. Chickens are slaughtered on-farm.	Chickens live continuously on pasture from the age of 4 weeks. This Step requires slower growing breeds that have the ability to perch. Chickens spend their entire lives on a single farm, and are slaughtered at a mobile or on-farm slaughter facility.

How to Read these Standards

Standards applicable to a Step level are designated with a ● symbol in the corresponding Step column. The ⓘ indicates the standard is considered a major non-conformance (see Non-conformances section below for more details, and the G.A.P. Policy Manual).

In the example below, the standard is required for each Step level, Step 1 through Step 5+:

STANDARD		Step Level					
		1	2	3	4	5	5+
2.4 Handling							
	Chickens must be handled properly and respectfully.	●	●	●	●	●	●
2.4.1 ⓘ	ⓘ [2.4.1 a]: G.A.P. has a zero-tolerance policy to kicking, throwing, striking, punching, hitting and any other actions or omissions in care that could cause harm or injury to chickens.						
	ⓘ [2.4.1 b]: Proper and respectful handling extends to all chickens – healthy, sick, injured and/or dead.						

In this example, 5.1.1 is required only for Step 3, 5.1.2 is required only for Step 4, and 5.1.3 is required for Step 5-5+. In addition, the ⓘ symbol prefaces additional information provided to aid in the understanding of the standard:

STANDARD		Step Level					
		1	2	3	4	5	5+
5.1 Outdoor and/or Pasture Access							
ⓘ Operations are strongly encouraged to provide chickens access to the outdoors from the youngest age possible and for as much of their lives as possible.							
5.1.1 ⓘ	All chickens from 4 weeks of age must be given continuous outdoor access during daylight hours unless climatic conditions pose a welfare risk.			●			

	<p>① [5.1.1]: Chickens in Step 3 systems have seasonal outdoor access.</p>						
5.1.2	<p>All chickens from 4 weeks of age must be given continuous access to pasture during daylight hours. If climatic conditions pose a welfare risk then chickens can be removed from pasture but must be given continuous access to an outdoor area during daylight hours. If the risk from climatic conditions is considered extreme, access to outdoor areas may be restricted as long as the restriction does not exceed 25 days (cumulative total) throughout a calendar year.</p> <p>① [5.1.2 a]: Chickens in Step 4 systems may only be confined to housing during extreme weather conditions (e.g. heavy precipitation, tornadoes, hurricanes, monsoons, blizzards, floods or non-typical weather for the season such as large swings in temperature) that jeopardize their welfare.</p> <p>① [5.1.2 b]: A calendar year is defined as January 1st through December 31st.</p> <p>① [5.1.2 c]: See Section 5.2 for requirements for pasture and outdoor areas.</p>					•	
5.1.3	<p>All chickens from 4 weeks of age must be given continuous access to pasture during daylight hours. If the risk from climatic conditions is considered extreme, access to outdoor areas may be restricted as long as the restriction does not exceed 5 consecutive days and 25 days (cumulative total) throughout a calendar year. Seasonal housing due to inclement climatic conditions is prohibited.</p> <p>① [5.1.3 a]: Chickens in Step 5 and Step 5+ systems may only be housed during extreme weather conditions (e.g. heavy precipitation, tornadoes, hurricanes, monsoons, blizzards, floods or non-typical weather for the season such as large swings in temperature) that jeopardize their welfare.</p> <p>① [5.1.3 b]: A calendar year is defined as January 1st through December 31st.</p> <p>① [5.1.3 c]: See Section 5.2 for requirements for pasture and outdoor areas.</p>					•	•

Program Requirements

The following is applicable to each operation applying for certification to Global Animal Partnership's 5-Step® Animal Welfare Standards for Chickens Raised for Meat.

1. General

- The standards in this document are requirements.
- The operation must have read the standards and prepared for the audit or they cannot be certified.
- With the exception of Standard 1.1.5 for Steps 5 and 5+, standards for breeding animals are not included in this document and will be developed in the future.
- Unless otherwise specified, standards in this document pertain to chickens of any age.
- The term "flock" is defined as a barn/house of chickens. In the event of split slaughter ages – for example if the flock is thinned – lameness and foot pad dermatitis assessments must be carried out in the oldest birds to be slaughtered.

- i. An example of a single flock could be one house of 10,000 chicks that are brooded together and then reared through to slaughter all on the same date.
 - ii. If chicks are brooded together but are subsequently raised to different Step levels, they would be considered two flocks and would require two sets of records.
 - iii. If 10,000 chicks are brooded together then split into two separate houses for this would be two flocks.
 - iv. If an operation has a single barn with two floors and the birds on the top floor are placed on one day and the birds on the bottom floor are placed 5 days later this would be two flocks.
- f. The term “operation” is defined as either (i) a single farm or (ii) a farm with more than one location, that meets all of the following criteria:
- i. all staff and chickens are under the direct supervision* of the main farm;
 - ii. the main farm owns all of the chickens; and
 - iii. the main farm owns, rents or leases, all the land and/or buildings where the chickens are raised.

An individual operation can include a farm that is under contract to raise chickens for a larger business OR a farm that owns their own chickens and markets them under their own brand(s).

**Direct supervision is defined as being when an employee of the main certified farm business is responsible for the chickens on the farm that is at a separate location to the main farm. If the person responsible for the management and care of the chickens at a separate location to the main certified farm business is a contract farmer and not an employee, then that site is a separate operation for the purposes of determining the number of audits – even if there is routine oversight from an employee of the main operation.*

- g. In order to achieve certification to a particular Step-level, the operation must meet all applicable standards. For example, all standards specified for Step 3 must be met in order to become certified to Step 3.
- h. The G.A.P. Policy Manual is a companion document to the standards, and details additional program requirements and terms of certification beyond that which is included in the standards (see www.globalanimalpartnership.org for a copy of the Policy Manual).
- i. G.A.P. is developing a Chain of Custody Records Reconciliation Program (CCRRP) and when finalized, all supply chains identified in the CCRRP will be required to be certified. CCRRP certification will include verifying numbers of animals/raw materials through the entire supply chain (i.e. from hatch to market).
- j. Each operation must follow a chain of custody program (see Section 9.6) that is also maintained by the slaughter facility and any further processor. The chain of custody program can be developed and implemented by an affiliated group (e.g., a producer group, co-operative, marketing entity) or created with the aid of external consultation.
- k. A glossary defining specific terms and terminology used in these standards is located at the end of this document.
- l. No standard in this document supersedes governmental regulations or laws, whether local, regional, state, provincial, territorial, federal, national, or other.

2. Applications

- a. Each operation is required to submit a completed G.A.P.’s 5-Step® Animal Welfare Chicken application for each certification cycle prior to audit.
- b. Each operation must identify all sites (either owned, leased, and/or shared) used to raise meat chickens by the operation on their G.A.P. application.
- c. Applications, this document, and other program documents, can be downloaded at www.globalanimalpartnership.org/ or by contacting your G.A.P.-accredited certifier.

3. Audit and Certification


- a. Each operation must submit a completed G.A.P.'s 5-Step® Animal Welfare Chicken application and be audited and certified prior to marketing any product(s) as G.A.P. Certified.
- b. Each operation must be audited once every certification cycle. A certification cycle is 15 months, which allows for chickens and operations to be assessed seasonally over a 5-year period.
- c. Each operation must have chickens on-site at the time of on-site audit, but not all houses at an operation must have chickens in them at the time of the on-site audit.
- d. At initial audit, the operation must have chickens on-site that are over 21 days old.
- e. At the time of on-site audit, the person(s) responsible for managing the operation and/or an animal caretaker must be present. A designated representative affiliated with a supplier group may also be present at the time of the on-site audit, but cannot be the only person present.
- f. Each operation applying for G.A.P. certification is responsible for ensuring that all required records and documents are available, and that all applicable standards are met, including actions that may be contracted or managed by another entity (e.g. catching and loading, transport, rodent control).
- g. All applicable standards, including those that may be controlled or managed by, or contracted to, another (e.g. the hatchery; a catching and loading crew; a transporter; a producer group, co-operative, or marketing entity; slaughter facility), will be assessed for compliance by the certification company and incorporated into its overall assessment of the operation prior to the final Step determination.
- h. G.A.P. supports the use of video or other electronic monitoring records for the review of brooding (for individual operations that market their own chickens, see n. ii. below) and catching and loading chickens. Use of video technology is not a requirement but can be used in place of certain observations listed in G.A.P.'s Policy Manual. Please refer to G.A.P.'s Policy Manual for additional details about how this must be conducted.
- i. Auditors do not make Step-level determinations nor provide consultative service to producers on meeting standards requirements; reviewers of authorized certification companies make Step-level determinations.
- j. Operations are only considered seasonal if they produce chickens for a portion of the year (e.g. flocks raised from July to November). If an operation produces chickens year-round, but only produces G.A.P. Certified chickens seasonally (e.g. for the holidays), the operation is not considered a seasonal operation.
- k. For operations that (1) only raise chickens seasonally, or (2) are new operations, the issue date of the G.A.P. certificate will be the date of the on-site audit, rather than the date the final Step-level decision is made.
- l. If chickens are not raised on a single operation and different stages of production are conducted by different operations (e.g. chickens are brooded and raised to 4 weeks at location A, and then transported to location B where from 4 weeks onwards they are raised to slaughter age), each operation must submit a completed G.A.P.'s 5-Step® Animal Welfare Chicken application and be audited and certified prior to product being marketed as G.A.P. Certified.
- m. Step-wise differentiation cannot be awarded on the basis of season. For example, a Step 3 operation cannot market their chickens as Step 3 during the time of year when chickens would typically be outdoors, and as a Step 2 when chickens would typically be kept indoors.
- n. **Specifics for Individual Operations that Market Their Own Chickens**
 - i. An operation that markets its own chickens must have at least 1 flock inspected when chickens are within 14 days of slaughter.
 - ii. At least 1 flock must be observed during brooding (up to 21 days) (this can be observed using video technology – see 3 h. above; G.A.P.'s Policy Manual; and check and confirm certifier's protocols for further details).
 - iii. If the operation has more than 4 flocks, then a minimum of 50% of the flocks will be selected by the auditor for inspection while on-site with preference given to the oldest flocks. If the operation has multiple locations, the 50% requirement must include at least one inspection at each location.




- iv. Catching and loading must be observed and audited at least once each certification cycle (this can be observed using video technology – see 3 h. above; G.A.P.’s Policy Manual; and check and confirm certifier’s protocols for further details).
 - v. The catching and loading observation referenced above can be of non-G.A.P. Certified chickens, provided all aspects of the catching and loading process are the same for all flocks, including those to be marketed as G.A.P. Certified.
- o. Specifics for Operations that are Part of an Affiliated Group**
- i. For operations that market chickens through an affiliated group (e.g. producer group, co-operative, marketing entity) rather than under the operation’s own name or brand, 30% of the operations must have 1 flock inspected when chickens are within 14 days of slaughter; 5% at less than 21 days of age; and the remaining 65% when the chickens are over 21 days of age.
 - ii. If an individual operation within an affiliated group has more than 4 flocks, then a minimum of 50% of the flocks will be selected by the auditor for inspection while on-site. If the operation has multiple locations, the 50% requirement must include at least one inspection at each location.
 - iii. Catching and loading must be observed and audited at least once each certification cycle. If the affiliated group controls catching and loading for multiple operations, the auditor may observe and inspect the catching and loading process for the group rather than on each applicant’s own operation, provided that (1) the assessment is performed at least once per certification cycle, and (2) all aspects of the catching and loading process are the same for all operations. (This can be observed using video technology – see 3 a. vii. Above; G.A.P.’s Policy Manual; and check and confirm certifier’s protocols for further details).
 - iv. The catching and loading observation referenced above can be of non-G.A.P. Certified chickens, provided all aspects of the catching and loading process are the same for all flocks, including those to be marketed as G.A.P. Certified.

4. Non-conformances

Note: This section provides a brief overview of the provisions of the G.A.P. Policy Manual relating to non-conformances. For further details please refer to that document.

- a. If an operation fails to meet a standard, it will be considered a non-conformance. There are three categories of non-conformance: minor, major and critical (see G.A.P.’s Policy Manual).
- b. If an operation receives a repeat non-conformance at the time of the next audit – the designation of minor, major and critical impacts the certification decision (see Repeat Non-conformances in G.A.P.’s Policy Manual).

In the example below, the  indicates that failure to meet the standard would be considered a critical non-conformance and the operation would be denied certification (see G.A.P.’s Policy Manual for further information).

STANDARD		Step Level					
		1	2	3	4	5	5+
2.4 Handling							
	Chickens must be handled properly and respectfully.	•	•	•	•	•	•
2.4.1 	<p> [2.4.1 a]: G.A.P. has a zero-tolerance policy to kicking, throwing, striking, punching, hitting and any other actions or omissions in care that could cause harm or injury to chickens.</p> <p> [2.4.1 b]: Proper and respectful handling extends to all chickens – healthy, sick, injured and/or dead.</p>						

In the example below, the **M** indicates that failure to meet the standard would be considered a major non-conformance. If at recertification, this standard was still not in compliance, then it would be considered a critical non-conformance and the operation would be denied certification (see G.A.P.'s Policy Manual for further information).

STANDARD		Step Level					
		1	2	3	4	5	5+
1.1 Source / Breed / Lines							
1.1.1 M	Intentional use of genetically modified or cloned chickens, or their progeny, is prohibited.	•	•	•	•	•	•

In the example below, as there is no **M** or **C** beside the standard, this means that the standard is classified as a minor non-conformance if it fails to be met. If at recertification, this standard was still not in compliance, then it would be considered a major non-conformance (see G.A.P.'s Policy Manual for further information).

STANDARD		Step Level					
		1	2	3	4	5	5+
2.2 Hazard Management							
2.2.1	Equipment, fittings, openings, protrusions, housing, outdoor areas, fences, and any other structures must be maintained to prevent injury.	•	•	•	•	•	•

- c. If an operation is issued a critical non-conformance, certification will be denied.
- d. If an operation is issued a major or minor non-conformance, it needs to be addressed and the response submitted to the certifier by the operation within 3 weeks from the date the certifier issues the audit report, and be accepted by the certifier as an acceptable response to the non-conformance, before a certificate is issued. If the operation does not respond to the non-conformance within 3 weeks, it will result in a shortened certificate once an acceptable response is received (see G.A.P.'s Policy Manual).

5. Step Differentiation within the Standards

- a. Each Step level—Step 1 through Step 5+—has its own requirements that must be met to be certified to that level. If an operation, for example, meets some, but not all Step 4 (or higher) requirements, but 100% of the requirements for Step 1, the G.A.P. certification will be Step 1.
- b. Step differentiator standards are those that do not apply to all steps. Meat chickens can achieve Step-levels 1, 2, 3, 4, 5 and 5+. Any standard that does not apply to all of those levels is a Step differentiator standard. In the example below Standard 4.5.4 only applies to Steps 1 and 2, Standard 4.5.5 only applies to Steps 3 and 4, and Standard 4.5.6 only applies to Steps 5 and 5+.

STANDARD		Step Level					
		1	2	3	4	5	5+
4.5 Lighting							
4.5.4 M	By day 3 after placement (except when raised under natural lighting conditions and dark periods are shorter), chickens must be provided daily with at least 8 hours of continuous light and at least 6 hours of continuous darkness (<1 lux) throughout their lives.	•	•				

4.5.5	By day 3 after placement (except when raised under natural lighting conditions and dark periods are shorter), chickens must be provided daily with at least 8 hours of continuous light and at least 8 hours of continuous darkness (<1 lux) throughout their lives.			•	•		
4.5.6	From placement (except when raised under natural lighting conditions and dark periods are shorter), chickens must be provided with a daily minimum of 8 hours of continuous darkness throughout their lives.				•	•	

- c. At renewal, if an operation fails to meet a standard that is specific to their level, it will drop to the applicable Step level, or lose certification as appropriate, unless the Certifier applies their discretion to issue a non-conformance **if and only if**:
- the standard that is out of conformance is not a repeat from the previous audit; and
 - the Certifier is confident the operation will be able to achieve and maintain the level specified in the standard; and
 - the operation is only out of conformance with one standard relating to the Step level the operation is looking to achieve.

6. Step-levels

- Step-level certification information (see Standard 9.6.2) must travel with chickens whenever they are being moved off the operation if they are to be marketed as G.A.P. Certified.
- All locations must carry the same Step-level or the lowest Step-level is applied to the marketed product.
 - For example, suppose there are five operations owned by different operators that produce meat chickens marketed under a single brand and chickens are not segregated according to operation. Prior to marketing chickens as G.A.P. Certified, all 5 operations need to be audited. Of these five operations four are certified to Step 4 and one is certified to Step 3. Because chickens are not segregated by operation, all chickens from all five operations would have to be marketed as Step 3.
 - As a second example, suppose there are six operations owned by different operators that raise chickens marketed by a single producer group with chickens segregated according to operation. Prior to marketing product as G.A.P. Certified, all six operations need to be audited. Of those six operations, four are certified as Step 4 and two are certified as Step 1. As the chickens are segregated according to operation, the chickens from the Step 1 operations can be marketed as Step 1 and the chickens from the Step 4 operations can be marketed as Step 4.
 - If chickens to be marketed as G.A.P. Certified are not raised on a single operation and different stages of production are conducted by different operations, each operation must submit a completed G.A.P.'s 5-Step[®] Animal Welfare Chicken application and be audited and certified, and the lowest Step-level achieved will be assigned to the marketed product. For example, chicks are brooded at Farm A, which sells the chickens to Farm B for grow-out. Farm A achieves a Step-level of Step 2, and Farm B is certified to Step 3. The final product will be labeled as Step 2.
- If an operation produces both G.A.P. Certified chickens and non-G.A.P. Certified chickens on the same site (e.g. where there are multiple houses managed to different programs), this is defined as a split operation. In order to qualify as a split operation, a strict segregation protocol must be in place and approved by the certifier prior to a G.A.P. certification decision being issued. The segregation protocol must include all of the following components (see G.A.P.'s Policy Manual):
 - A written policy describing how chicken from G.A.P. Certified flocks is segregated from chicken from non-G.A.P. Certified flocks; AND
 - How chickens are physically separated (for example packed on dedicated lines); OR
 - How chickens are identified using a method that allows for instant visual identification.
- As outlined in the G.A.P. Policy Manual, split operations will not be issued certification to the operation as a whole, rather certificates will specify the certified G.A.P. Certified portion of the split operation.

7. Additional Standards Documents

Note: The most current versions of the documents listed below can be downloaded at www.globalanimalpartnership.org.

- a. As noted above, G.A.P.'s 5-Step® Animal Welfare Standards for Chickens Raised for Meat v3.2 is accompanied by G.A.P.'s Policy Manual.
- b. As per bullet 2a above, each operation is required to complete G.A.P.'s 5-Step® Animal Welfare Chicken application each certification cycle.

8. Labeling

- a. Prior to labeling any chicken as G.A.P. Certified, the marketing entity must ensure that they meet labeling regulations for the country in which they are selling.
- b. The use of the G.A.P. label/logo must meet the requirements of G.A.P.'s Labeling Guide, and be approved by G.A.P. prior to use.
- c. Before retail-ready packaged products are labeled as G.A.P. Certified, operations must contact G.A.P.'s Labeled Products Authorization Program at lpa@globalanimalpartnership.org for details of the application, fee schedule and approval process.

TABLE OF CONTENTS


1 ANIMAL SOURCE AND HEALTH					
Source / Breed / Lines	12	Litter and Flooring	21	Guardian Animals	34
Growth Rates	13	Air Quality	22	9 PLANS, PROTOCOLS, PROCEDURES, TRAINING, RECORDS AND DOCUMENTS	
Medication	13	Lighting	22	General Requirements	35
Treatment	13	Stocking Density	23	Written Farm/Animal Health Plan/ System Plan	35
On-Farm Euthanasia	14	Perches	24	Biosecurity Procedures and Protocols	36
		Environmental Enrichments	24	Alternative Power Supply	36
2 ANIMAL CARE AND MANAGEMENT		5 OUTDOOR CONDITIONS		Training	36
Daily Flock Management	16	Outdoor and/or Pasture Access	26	Traceability and Chain of Custody	37
Hazard Management	16	Vegetation and Foraging Materials	27	10 SLAUGHTER REQUIREMENTS	
Chicks and Brooding	16	Space Requirements of Outdoor Areas	27	General Requirements	38
Handling	16	Provisions to Encourage Chickens to Use Outdoor and Pasture Areas	28	Segregation and In-Plant Traceability	39
Physical Alterations	16	6 LOADING		Appendix I: Breeds/Strains Growth Rates and Step-levels	40
Thermal Comfort	17	Condition of Chickens	29	Appendix II: Identifying and Scoring Footpad Dermatitis	42
Feather Pecking	17	Water and Feed Withdrawal	29	Appendix III: Intervention Plans	45
Footpad Dermatitis	17	Catching and Loading	29	Appendix IV: 3-Category Lameness Evaluation	46
Lameness	18	Transport Containers	30	Appendix V: Soiled Feather Assessment	48
Mortality	18	7 TRANSPORT		Appendix VI: Litter Quality Assessment	50
Culling	19	Transport of Chicks	31	Appendix VII: Evaluation of Air Quality	52
Other Commercially Raised Animals on the Operation	19	Equipment and Vehicles for Chickens	31	Appendix VIII: Environmental Enrichments	54
Domestic Animals on the Operation	19	Transport Personnel Responsibilities and Procedures	31	Appendix IX: Illustration of “At Chicken Height”	71
3 FEED AND WATER		Transport Duration for Chickens	32	Appendix X: Provisions to Encourage Chickens to Use Outdoor and Pasture Areas	72
Water Availability	20	Transport Records	32	Glossary	75
Feeding Requirements	20	8 RODENT, WILD BIRD, AND PREDATOR CONTROL			
Feed Hygiene	20	Rodent Control	33		
Additives and Ingredients in Feed or Water	20	Wild Bird Control	33		
4 HOUSING		Predator Control	33		
Housing Systems	21				
Access to Housing	21				



① Certain historical records and documents included in this set of standards may not be available at the time of initial audit as the operation applying for 5-Step® certification was unaware they would be required to monitor and/or record them and, therefore, cannot create them for past events, treatments, assessments, or practices. At the time of initial audit, record-keeping, protocols and documentation mechanisms must be in place to meet each of these standards, and be available for review.

1 ANIMAL SOURCE AND HEALTH

① G.A.P. announced in March 2016 that it would be transitioning all of our Step Levels to higher welfare genetics by 2024. To assist us to that end, G.A.P. has provided a research grant to the University of Guelph to determine the specific requirements for each Step level. Once the findings from that study are available, G.A.P. will use the information to update this document, and publish as v4.0 of this standard.

① It is G.A.P.'s intention to develop standards for parent stock (i.e. broiler breeders) as a component of v4.0.

STANDARD	Step Level					
	1	2	3	4	5	5+
1.1 Source / Breed / Lines ① The purpose of this Section is to ensure that producers choose the right breed or strain for the system they are managing, whether this is indoors or pasture based. The over-arching principles for good broiler chicken welfare that must be considered when selecting a breed or strain are covered in Standards 1.1.2 to 1.1.4. The informational notes below Standards 1.1.2 to 1.1.4 reference related sections within the standard that can be used to substantiate breed or strain suitability.						
1.1.1 	Intentional use of genetically modified or cloned chickens is prohibited.					
1.1.2	Breeds/lines/strains must be chosen for good leg health and for low levels of mortality.					
	① [1.1.2]: See related Sections 2.9 and 2.10.					
1.1.3	Breeds/lines/strains must be chosen for the ability to range and for good immune systems.					
	① [1.1.3]: See related Sections 1.4.2, 2.10, 5.1, 5.2, 5.3, 5.4.					
1.1.4	Breeds/lines must be able to perch throughout their lives.					
	① [1.1.4 a]: G.A.P.'s 5-Step® Animal Welfare Standard defines perching as the act of balancing with feet (claws) wrapped around an elevated object such as a wooden bar or pole, or a tree branch. Chickens must be able to perch while in both sitting and standing positions.					
	① [1.1.4 b]: See related Section 4.7 and Standard 5.4.3.					
1.1.5	Each operation must have a signed affidavit from the source confirming that parent stock (i.e. broiler breeders) meet <u>all</u> of the following criteria:					
	a. fed daily; b. not water restricted; c. never been kept in cages; and d. never given sub-therapeutic antibiotics.					
	① [1.1.5]: The affidavit may come from the hatchery or the breed company.					

STANDARD		Step level						
		1	2	3	4	5	5+	
1.2 Growth Rates								
<p>① Producers are not required to weigh their chickens – compliance is currently determined by G.A.P. from the breeder’s published data.</p> <p>① See Appendix I for a list of breeds/strains, their average growth rate determined from the breeder’s published data, and the maximum step level they can achieve. Any operation with questions around acceptable breeds/strains should contact G.A.P. directly prior to scheduling their audit.</p>								
1.2.1	The maximum average growth rate must not exceed 68 grams (0.150 lbs) per day.	•	•	•				
1.2.2	The maximum average growth rate must not exceed 50 grams (0.110 lbs) per day.				•			
1.2.3	The maximum average growth rate must not exceed 45 grams (0.099 lbs) per day.					•		
1.2.4	The maximum average growth rate must not exceed 35 grams (0.077 lbs) per day.						•	
1.3 Medication								
1.3.1 	Chickens that are given antibiotics, ionophores, beta agonists, sulfa drugs and/or arsenic-based drugs are prohibited from being marketed as G.A.P. Certified.	•	•	•	•	•	•	
	<p>① [1.3.1 a]: This standard applies whether treatments are given therapeutically or sub-therapeutically.</p> <p>① [1.3.1 b]: See Standard 1.4.1 for prompt treatment of chickens.</p> <p>① [1.3.1 c]: Arsenic-based drugs include, but are not limited to 3-Nitro®, Roxarsone, Nitarsone, Arsanilic Acid, and Carbarsonne.</p>							
	1.3.2	A written protocol must be in place to identify and ensure that any chickens treated with antibiotics, ionophores, beta agonists, sulfa drugs and/or arsenic-based drugs are not marketed as G.A.P. Certified.	•	•	•	•	•	•
	1.3.3	Off-label / extra-label use of medicines is prohibited unless prescribed or advised by a veterinarian.	•	•	•	•	•	•
<p>① [1.3.3 a]: Veterinarian prescription documentation may be acquired via email or fax.</p> <p>① [1.3.3 b]: Parasiticides including coccidiostats and vaccines are not included in this standard.</p>								
1.3.4	Expired medication is prohibited.	•	•	•	•	•	•	
1.4 Treatment								
1.4.1 	Sick or injured chickens must be promptly treated or euthanized according to Section 1.5.	•	•	•	•	•	•	
1.4.2	Records must be kept of <u>any</u> treatment (medication, vaccinations, probiotics etc.) to any individual or group of chickens including:							
	<ul style="list-style-type: none"> a. any substance administered; b. date and method of administration; and c. flock/bird ID. 	•	•	•	•	•	•	
1.4.3	Veterinarian-prescribed treatments must be administered according to veterinarian guidance.	•	•	•	•	•	•	
1.4.4	Any area designated for sick or injured chickens, such as a hospital pen or designated area within an existing pen, must meet the space requirements and housing conditions detailed in the HOUSING section.	•	•	•	•	•	•	
	<p>① [1.4.4]: Sick and/or injured chickens may be segregated from healthy chickens when necessary, but it is not required if it is in the best interest of the bird to keep it with the flock.</p>							


STANDARD		Step level					
		1	2	3	4	5	5+
1.4 Treatment Continued							
1.4.5	Segregated sick or injured chickens must be monitored at least twice daily.	●	●	●	●	●	●
1.4.6	Operations must have an internal and external parasite control program that can be implemented if parasites are impacting bird health and welfare.	●	●	●	●	●	●
1.4.7	Products containing organophosphates, cannot be applied directly to chickens.	●	●	●	●	●	●
1.5 On-Farm Euthanasia							
<p>① This Section and the methods listed in Standard 1.5.7 relate to on-farm euthanasia, which G.A.P. defines as the act of killing individual chickens in response to an irrecoverable illness or injury.</p> <p>① If emergency slaughter of the whole flock or a large proportion of the flock needs to take place due to a disease outbreak, this Section does not restrict the actions of the farm if the conditions or circumstances require them to take action guided by a veterinarian or regulatory official.</p>							
1.5.1	Euthanasia must be performed by a trained person or veterinarian.	●	●	●	●	●	●
	① [1.5.1]: Producers will not be required to demonstrate the ability to euthanize a bird in order to show compliance with this standard, but they must be able to describe the training they received whether experiential or formal.						
1.5.2	Any bird identified as requiring euthanasia must be euthanized within 4 hours of discovery.	●	●	●	●	●	●
	① [1.5.2]: Timely euthanasia is vital. Ideally any bird identified as requiring euthanasia will be euthanized immediately; however, G.A.P. understands that it may take time for a trained person or the correct equipment to get to the operation.						
1.5.3	Euthanasia technique(s) must cause rapid and irreversible insensibility immediately followed by death.	●	●	●	●	●	●
	① [1.5.3]: The operation must be able to articulate to the auditor the visual indicators of death, and the physical parameters that confirm this.						
1.5.4	The person performing euthanasia must remain with the chicken(s) until death is evident.	●	●	●	●	●	●
1.5.5	Euthanasia equipment must be maintained according to manufacturer's specifications.	●	●	●	●	●	●
1.5.6	All euthanized/dead chickens must be removed immediately from housing and/or outdoor areas and pasture in use.	●	●	●	●	●	●
	① [1.5.6]: It is the responsibility of the operation to dispose of dead chickens according to local, state, provincial, territorial, and/or federal or national regulations. Removal can include burial or composting in designated areas that will not put chickens at risk from transmission of diseases and/or the attraction of predators.						


STANDARD		Step level										
		1	2	3	4	5	5+					
1.5 On-Farm Euthanasia <i>Continued</i>												
1.5.7	Methods of euthanasia are listed below, where YES indicates an acceptable method and NO indicates an unacceptable method. Chickens must be appropriately held or restrained as necessary to ensure the euthanasia method can be properly and safely administered.											
	METHOD	ACCEPTABILITY										
	Manual cervical dislocation (i.e. use of hands only to dislocate the neck as near to the head or skull as possible)	YES										
	Penetrating captive bolt pistol ¹	YES										
	Non-penetrating captive bolt pistol ¹	YES										
	Electrical stun knife ¹ (only permitted if chickens are stunned prior to cutting the neck)	YES										
	Gas stunning and killing systems ¹ using (1) multi-phase carbon dioxide ² , (2) argon, (3) nitrogen, or (4) a mixture of these gases	YES										
	Veterinarian administered overdose of injectable anesthetics, including barbiturate and barbituric acid derivatives	YES					•	•	•	•	•	•
	Mechanical cervical dislocation (i.e. equipment that pulls/crushes the neck such as wringers or poultry pliers or handheld cervical dislocators such as the Koechner Euthanasia Device)	NO										
	Manually applied blunt force trauma to the head	NO										
	Decapitation	NO										
	Bleeding/slitting the throat without pre-stunning	NO										
	De-braining (inserting a sharp implement through the roof of the chicken's mouth into its brain)	NO										
	Gas stunning and killing systems using carbon monoxide	NO										
¹ Only permitted if used to the manufacturer's specifications.												
² Mult-phase carbon dioxide systems must have at least two phases where the first phase has a lower concentration of carbon dioxide to render the chickens unconscious before higher levels of carbon dioxide are introduced.												
<p>i [1.5.7 a]: If an operation uses a method of euthanasia not listed above, written approval from Global Animal Partnership must be received prior to on-farm use in order to meet this Standard.</p> <p>i [1.5.7 b]: Fatigue can be an issue for caretakers when multiple chickens require euthanasia, so G.A.P. encourages appropriate staffing and training.</p>												

2 ANIMAL CARE AND MANAGEMENT












STANDARD		Step Level					
		1	2	3	4	5	5+
2.1 Daily Flock Management							
2.1.1	Each flock must be observed and monitored at least twice daily. Records of observation and monitoring must be kept.	•	•	•	•	•	•
	ⓘ [2.1.1 a]: Each inspection, whether twice daily or more, must be recorded to meet this standard. ⓘ [2.1.1 b]: See also standard 1.4.5 for the observation of individual chickens that have been segregated from the flock.						
2.2 Hazard Management							
2.2.1	Equipment, fittings, openings, protrusions, housing, outdoor areas, fences, and any other structures must be designed and maintained to prevent injury.	•	•	•	•	•	•
2.2.2	Chickens must be kept from contact with any potentially toxic or injurious substances (e.g., those used for maintenance, sanitation, cleaning, insect and rodent control).	•	•	•	•	•	•
2.3 Chicks and Brooding							
ⓘ See also Section 7.1.							
2.3.1	Placement of chicks must begin within 2 hours of delivery to the operation.	•	•	•	•	•	•
	ⓘ [2.3.1]: Chick delivery boxes should be kept low to the ground and tipped in a manner that does not cause injury during unloading.						
2.3.2	Houses must be pre-warmed prior to chick arrival to accommodate their thermal needs.	•	•	•	•	•	•
	ⓘ [2.3.2]: At chick height, a minimum temperature of 90°F (32°C) is recommended, but chick behavior should be the ultimate determinant of their thermal comfort.						
2.4 Handling							
2.4.1	Chickens must be handled properly and respectfully.	•	•	•	•	•	•
	ⓘ [2.4.1 a]: G.A.P. has a zero-tolerance policy to kicking, throwing, striking, punching, hitting and any other actions or omissions in care that could cause harm or injury to chickens.						
	ⓘ [2.4.1 b]: Proper and respectful handling extends to all chickens – healthy, sick, injured and/or dead.						
2.5 Physical Alterations							
2.5.1	All physical alterations are prohibited, including, but not limited to: <ol style="list-style-type: none"> beak trimming / beak conditioning; dubbing; caponization; de-spurring; and toe trimming/toe clipping/de-toeing/toe conditioning. 	•	•	•	•	•	•

STANDARD		Step Level					
		1	2	3	4	5	5+
2.6 Thermal Comfort							
2.6.1	The thermal comfort of chickens must be maintained at all times through management and the provision of supplemental heating and/or cooling, as necessary.	•	•	•	•	•	•
	📌 [2.6.1]: The auditor will assess whether chickens show signs of thermal discomfort at audit.						
2.7 Feather Pecking							
2.7.1	Any evidence of feather pecking must be promptly addressed and managed.	•	•	•	•	•	•
2.7.2	Records of any evidence of feather-pecking incidences are required, including:						
	<ul style="list-style-type: none"> a. date of incidence; b. percentage of affected chickens per flock (injured chickens, chickens with feather loss, chickens that die from injurious pecking); c. actions taken to address the incidence; and d. outcomes of the actions taken to address the incidence. 	•	•	•	•	•	•
	📌 [2.7.2]: Records are required only if incidence(s) occur.						
2.7.3	The use of goggles, blinkers, contact lenses, or any other artificial devices is prohibited.	•	•	•	•	•	•
2.8 Footpad Dermatitis							
📌 See Appendix II for footpad scoring. Note: the sums below are not based only on prevalence, but rather a calculation that accounts for both prevalence and severity of the footpad lesions.							
2.8.1 ^M	Each operation must have a footpad dermatitis monitoring program (per Appendix II) that is conducted either at the slaughter facility OR on-farm.	•	•	•	•	•	•
	📌 [2.8.1 a]: See Appendix II: Identifying and Scoring Footpad Dermatitis for charts of the scoring protocol.						
	📌 [2.8.1 b]: If chickens are scored on-farm, this must occur during the final week prior to slaughter.						
	📌 [2.8.1 c]: If scoring is carried out at the slaughter facility the same representative sample of birds must be scored as would be scored on-farm. See Appendix II for further details and examples.						
2.8.2	Records of footpad dermatitis and the corresponding footpad dermatitis sum from the monitoring sample must be kept for each flock.	•	•	•	•	•	•
	<ul style="list-style-type: none"> 📌 [2.8.2 a]: See Appendix II: Identifying and Scoring Footpad Dermatitis for specifications on calculating the flock's total footpad dermatitis sum. 📌 [2.8.2 b]: Only scores collected during the final week prior to slaughter can be used in determining the sum for each flock. 						
2.8.3	Footpad dermatitis sums must not exceed an annual average score of 20.	•					
2.8.4	Footpad dermatitis sums must not exceed an annual average score of 15.		•	•			
2.8.5	Footpad dermatitis sums must not exceed an annual average score of 5.				•	•	
2.8.6	Footpad dermatitis sums must not exceed an annual average score of 2.						•

STANDARD		Step level					
		1	2	3	4	5	5+
2.8 Footpad Dermatitis <i>Continued</i>							
2.8.7	If the incidence of footpad dermatitis exceeds the footpad dermatitis scores detailed above for any individual flock, an intervention plan, as detailed in Appendix III, that addresses, at a minimum, feed composition, litter management, humidity control, and stocking density must be implemented immediately to reduce levels in subsequent flocks.	•	•	•	•	•	•
	<p>① [2.8.7 a]: Operations may find it useful to check footpads earlier in the growing period, but any checks done prior to slaughter do not replace the requirements in 2.8.1.</p> <p>① [2.8.7 b]: An annual average score is used in 2.8.3 to 2.8.6 to accommodate seasonal variability in footpad dermatitis scores; however, operations are required to manage each individual flock that exceeds the relevant score.</p>						
2.9 Lameness							
① See Appendix IV for lameness evaluations. Note: the sums below are not based only on prevalence, but rather a calculation that accounts for both prevalence and severity of lameness.							
2.9.1 	Lameness evaluations must be conducted on the operation, as described in Appendix IV, during the final week prior to slaughter.	•	•	•	•	•	•
	① [2.9.1]: See Appendix IV: 3-Category Lameness Evaluation for scoring protocol, sample size, and specification on calculating the flock's total lameness sum.						
2.9.2	Lameness records for each flock and the corresponding lameness sum from the monitoring sample for each flock are required.	•	•	•	•	•	•
	① [2.9.2]: See Appendix IV: 3-Category Lameness Evaluation for specifications on calculating the flock's total lameness sum.						
2.9.3	Lameness score for each individual flock must not exceed 20.	•					
2.9.4	Lameness score for each individual flock must not exceed 15.		•	•			
2.9.5	Lameness score for each individual flock must not exceed 5.				•		
2.9.6	Lameness score for each individual flock must not exceed 2.					•	•
2.9.7	If the lameness evaluation conducted during the final week prior to slaughter exceeds the Step-differentiation total detailed above, a written intervention plan, as detailed in Appendix III, is required to reduce levels in subsequent flocks.	•	•	•	•	•	•
2.10 Mortality							
① Flock mortality (from placement) thresholds in Standards 2.10.1 – 2.10.6 do not include predated chickens or culls.							
① There are four categories of mortality: (a) chickens found dead through natural causes, (b) chickens found dead through predation, (c) chickens that are missing (chickens lost or suspected to be predated) and (d) chickens that are culled (see Standard 1.4.1).							
① Though all categories of mortality must be recorded, the percentages used to calculate the flock mortality thresholds in Standards 2.10.2 to 2.10.4 include only chickens that die through natural causes - point (a) above. Losses due to predation and/or lost or missing chickens are not included in this calculation.							
① Missing birds - point (c) above - are only recorded once for each flock, at removal for slaughter.							
2.10.1	Daily records that separately identify chickens that die, are culled and are predated are required for each flock.	•	•	•	•	•	•

STANDARD		Step level					
		1	2	3	4	5	5+
2.10 Mortality Continued							
2.10.2 	Annual average flock mortality must not exceed 6%.	•					
2.10.3	Annual average flock mortality must not exceed 5%.		•	•			
2.10.4	Annual average flock mortality must not exceed 4%.				•	•	•
2.10.5	If individual flock mortality exceeds the percentages above for any one flock throughout the year, a written intervention plan, as detailed in Appendix III, is required to reduce levels in the existing and subsequent flocks.	•	•	•	•	•	•
2.11 Culling							
2.11.1	Any chickens meeting the following criteria must be culled according to acceptable euthanasia methods and recorded: <ul style="list-style-type: none"> a. runts; b. exhibiting a lameness score of 2; and/or c. sick or injured without chance for recovery (for example sick or injured such that the chicken is unable to reach food and water). 	•	•	•	•	•	•
① [2.11.1]: See Appendix IV: 3-Category Lameness Evaluation for details on scoring lameness, and Section 1.5 for euthanasia requirements.							
2.12 Other Commercially Raised Animals on the Operation							
2.12.1	All species of animals raised commercially on the operation (for which G.A.P.'s 5-Step® Animal Welfare Standards exist) must be raised and handled according to Step 1 standards or higher.					•	•
① [2.12.1] Animals kept as pets, show animals or other non-commercial situations are excluded from this standard.							
2.13 Domestic Animals on the Operation							
① Domestic animals include dogs, cats, horses, or any other animals on the operation, including both pets and working animals.							
2.13.1	Neglect or abuse of domestic animals is prohibited.	•	•	•	•	•	•
2.13.2	All domestic animals on the operation must be provided with: <ul style="list-style-type: none"> a. food and water on a daily basis, as evidenced by healthy body condition score; b. surroundings that do not cause them injury; c. an environment that allows for freedom of movement and exercise; d. a comfortable resting area that provides protection from temperature extremes; and e. veterinary attention, if required. 					•	•

3 FEED AND WATER

STANDARD		Step Level					
		1	2	3	4	5	5+
3.1 Water Availability							
3.1.1 	All chickens must have continuous access to drinking water.	•	•	•	•	•	•
	 [3.1.1]: Water may be withheld for up to 2 hours for the purposes of then delivering supplements and/or medication such as vaccine(s) in water.						
3.1.2	Waterers must be checked daily and any debris cleaned out.	•	•	•	•	•	•
	 [3.1.2]: Cleaning out debris could include removal of dirt or waste matter from bell or pan waterers or flushing nipple lines (as required).						
3.2 Feeding Requirements							
3.2.1 	All chickens must have <i>ad-libitum</i> access to feed during daylight hours.	•	•	•	•	•	•
3.2.2	Feeders must be designed and distributed to allow chickens to eat without restriction.	•	•	•	•	•	•
3.2.3	Chickens fed whole grains and/or given pasture access must be provided with insoluble grit.	•	•	•	•	•	•
3.3 Feed Hygiene							
3.3.1	Feed in storage bins and feeders must not be moldy or mildewed, contaminated by rodents, or otherwise compromised in quality.	•	•	•	•	•	•
3.3.2	Feeders must be free of debris.	•	•	•	•	•	•
3.4 Additives and Ingredients in Feed or Water							
3.4.1 	Mammalian by-products are prohibited.	•	•	•	•	•	•
	 [3.4.1]: By-products include animal waste and products derived from slaughter/harvest process (see glossary).						
3.4.2 	Avian by-products, including eggs, are prohibited.	•	•	•	•	•	•
	 [3.4.2]: By-products include animal waste and products derived from slaughter/harvest process (see glossary).						
3.4.3 	Fish and fish by-products are prohibited.	•	•	•	•	•	•
	 [3.4.3]: This includes whole fish, parts of fish, fishmeal, fish by-products from the processing industry and other aquatic species and/or products (does not include seaweed or oyster shell).						
3.4.4	Each operation must keep up-to-date feed ration ingredient lists, or tags, including mineral/vitamin mixes whether using purchased or home mixed feed. Lists and tags need to be made available to the auditor.	•	•	•	•	•	•
	 [3.4.4]: If the operation is part of an affiliated group which determines the feed ration the group can provide the auditor with this potentially confidential and proprietary information.						

4 HOUSING

STANDARD	Step Level					
	1	2	3	4	5	5+
4.1 Housing Systems						
① <i>Movable pens/coops/huts without floors (also commonly known as tractors and arks) that confine chickens and only allow them to range within the unit can <u>only</u> achieve Step 2.</i>						
4.1.1 ①	All cage systems are prohibited.	•	•	•	•	•
	① [4.1.1 a]: A cage is a fully enclosed structure made of mesh, bars, or wires that prevents full range of motion and the ability to express natural behavior, such as roosting, foraging, and exercising (e.g., battery cage, caged broiler systems). Cages do not include fenced-in verandas and outdoor enclosures provided they allow for full range of motion and the ability to express natural behavior. Transport containers are also excluded from the definition of cages. ① [4.1.1 b]: This standard does not preclude the use of netted mobile/moveable coop systems, however these systems can <u>only</u> achieve Step 2.					
4.1.2	Group size must not exceed 2000 chickens.				•	•
	① [4.1.2]: A group is defined as a number of chickens associated with either i) a mobile house; ii) a single flock in a stationary house; or iii) a subdivision of chickens within a larger stationary house.					
4.2 Access to Housing						
4.2.1	All chickens must have continuous access to housing, whether mobile or permanent, that provides protection from the elements and predation.	•	•	•	•	•
4.3 Litter and Flooring						
4.3.1	Floors of all houses, including mobile houses that are stationary for more than 3 days, must be covered with a minimum total of 3 in (7.6 cm) of non-toxic litter.	•	•	•	•	•
4.3.2	Litter must be managed so it is friable and no more than 10% of the littered area is caked.	•	•	•	•	•
	① [4.3.2]: Caked litter areas directly under feed and water lines are included in the 10% allowable caked area.					
4.3.3	If litter is not completely removed (i.e. cleaned out) between flocks, all caked litter must be removed between flocks, and top dressed as necessary so that the entire floor is covered to a total minimum litter depth of 3 in (7.6 cm).	•	•	•	•	•
4.3.4	Litter must be of quality and quantity to:					
	a. provide a comfortable environment, and b. allow for dust-bathing behavior, and foraging/scratching.	•	•	•	•	•
4.3.5	A Soiled Feather Assessment (see Appendix V) must be conducted for each flock between 15-20 days of age. Records must be kept.	•	•	•	•	•
	① [4.3.5]: The operation must conduct the assessments as required by the standards. At the time of audit, auditors will perform an assessment of soiled feathers as an additional indicator of litter conditions (NB: the birds assessed by the auditor will not necessarily be 15-20 days of age). See Appendix V: Soiled Feather Assessment.					

STANDARD		Step Level					
		1	2	3	4	5	5+
4.3.6	A Litter Quality Assessment (see Appendix VI) must be conducted for each flock between 15-20 days of age. Records must be kept.	•	•	•	•	•	•
	① [4.3.6]: The operation must conduct the assessments as required by the standard. At the time of audit, auditors will additionally perform a Litter Quality Assessment. (NB: the birds assessed by the auditor will not necessarily be 15-20 days of age).						
4.3.7	If the assessment detailed in Standard 4.3.5 shows that more than 10% of sampled birds are severely soiled OR the litter score detailed in Standard 4.3.6 is more than 1, a written intervention plan, as detailed in Appendix III, designed to improve litter quality and ensure that feather soiling does not deteriorate further must be implemented that same day.	•	•	•	•	•	•
	① [4.3.7]: Litter quality and feather cleanliness are linked, therefore if either or both measures are outside the limits set by this standard an intervention plan to remediate the situation must be put into place.						
4.3.8	If the assessments detailed in Standards 4.3.5 and 4.3.6 show that more than 10% of sampled birds are severely soiled OR the litter score is more than 1, an additional Soiled Feather Assessment must be conducted for the same flock a week prior to slaughter. Records must be kept.	•	•	•	•	•	•
4.3.9	If the assessments detailed in standards 4.3.5 and 4.3.6 show that more than 10% of sampled birds are severely soiled OR the litter score is more than 1, an additional Litter Quality Assessment must be conducted for the same flock a week prior to slaughter. Records must be kept.	•	•	•	•	•	•
4.3.10	Slatted and/or wire floors are prohibited.	•	•	•	•	•	•
4.4 Air Quality							
① See Appendix VII: Sensory Evaluation of Air Quality protocol.							
4.4.1	If chickens are shut in housing or structures for any part of the day, air quality must be assessed at least once each day, using calibrated meters or testing strips, or sensory evaluation and records must be kept.	•	•	•	•	•	•
	① [4.4.1]: For systems where chickens have access to the outdoors/pasture air quality must be assessed before popholes are opened.						
4.4.2	Air quality measures must not exceed the following levels when calibrated meters or testing strips are used: a. dust: 10 mg per cubic meter; b. ammonia: 20 ppm;	•	•	•	•	•	•
	OR score more than 1 on the air quality scale in Appendix VII.						
4.4.3	If air quality is found to exceed the levels in Standard 4.4.2, a written intervention plan, as detailed in Appendix III, designed to improve air quality must be implemented that same day.	•	•	•	•	•	•
4.5 Lighting							
4.5.1	Light intensity in housing during daylight hours, either from artificial or natural light or a combination of artificial and natural light, must be at least 50 lux (4.6 foot candles).	•	•	•	•	•	•
	① [4.5.1]: Operations that meet Standards 4.5.2 and 4.5.5 (if applicable) do not additionally have to meet this standard.						

STANDARD		Step level					
		1	2	3	4	5	5+
4.5 Lighting Continued							
4.5.2	By 1 January 2022, all operations with chickens housed continuously, seasonally, or during brooding, must be provided with natural light through insulated windows, solar tubes or semi-transparent roofing placed evenly throughout barns/houses so that the combined area through which natural light enters is equal or greater than 1.0% of the total floor space. ① [4.5.2]: Operations utilizing curtain barns meet this standard providing the curtain area is equal or greater than 1.0% of the total floor space. See also Standard 4.5.4.		•	•	•	•	•
4.5.3	By 30 September 2018, operations must have a written transition plan detailing the changes they will make to meet Standard 4.5.2 by 1 January 2022.		•	•	•	•	•
4.5.4	From 1 October 2018, certified operations must start to implement the plan detailed in Standard 4.5.3 and begin the alterations necessary to transition barns to meet Standard 4.5.2. ① [4.5.4]: It is recommended that all operations start work to transition barns to provide natural light as soon as possible.		•	•	•	•	•
4.5.5	By 1 January 2022, when curtain barns are used to house chickens there must be a method of providing natural light to meet the requirement of Standard 4.5.2 if the curtains block the light completely when they are closed. ① [4.5.5 a]: This could be through having windows or solar tubes in the roof of the curtain barn, areas of semi-transparent roofing or use of translucent curtains. ① [4.5.5 b]: Additional natural light need not be provided if curtains are closed for no more than seven consecutive days per flock for the purposes of acclimatizing birds to cold weather in the winter.		•	•	•	•	•
4.5.6 Ⓜ	By day 3 after placement (except when raised under natural lighting conditions and dark periods are shorter), chickens must be provided daily with at least 8 hours of continuous light and at least 6 hours of continuous darkness (<1 lux) throughout their lives.	•	•				
4.5.7	By day 3 after placement (except when raised under natural lighting conditions and dark periods are shorter), chickens must be provided daily with at least 8 hours of continuous light and at least 8 hours of continuous darkness (<1 lux) throughout their lives.			•	•		
4.5.8	From placement (except when raised under natural lighting conditions and dark periods are shorter), chickens must be provided with a daily minimum of 8 hours of continuous darkness throughout their lives.					•	•
4.6 Stocking Density							
① Stocking density must be calculated using the final target weight of all chickens per flock per area at the time of catching plus a maximum of 4% variation to calculate the number of chickens to be placed within a given barn/house. Expected liveability must not be included in the calculation.							
4.6.1	Chickens must have enough space to express natural behavior, including standing, turning around, and preening, without touching another chicken.	•	•	•	•	•	•
4.6.2 Ⓜ	For birds placed from 1 January 2018 to 30 June 2020, stocking density must not exceed 6.5 lbs/ft ² (32 kg/m ²).	•	•				
	For birds placed from 1 July 2020 onwards, stocking density must not exceed 6 lbs/ft ² (29 kg/m ²).	•	•				
	① [4.6.2]: G.A.P. recognizes that Step 1 and 2 operations need time to adjust bird numbers, egg sets and potentially build new housing in order to meet the stocking density required from 1 July 2020 onwards. Operations should use the time allotted to make progress towards meeting the lower stocking density across all their Step 1 and Step 2 housing.						

STANDARD		Step Level					
		1	2	3	4	5	5+
4.6 Stocking Density <i>Continued</i>							
4.6.3	Stocking density must not exceed 6 lbs/ft ² (29 kg/m ²).			•			
4.6.4	Stocking density must not exceed 5.5 lbs/ft ² (27 kg/m ²). ⓘ [4.6.4 a]: This stocking density only applies if chickens are confined in housing without access to the outdoors. ⓘ [4.6.4 b]: If confinement of the flock needs to take place due to a disease outbreak, this Standard does not restrict the actions of the farm if the conditions or circumstances require them to take action guided by a veterinarian or regulatory official.				•	•	•
4.7 Perches							
4.7.1	Perches must be provided wherever birds roost – whether this is in the housing structure or outside on pasture - at a minimum of 7 in (18 cm) of perch space per bird to enable all chickens to perch simultaneously. ⓘ [4.7.1]: Perches are to be elevated off the floor, be easily used by the chickens, and positioned at heights to allow for use by chickens from an early age throughout their lives.					•	•
4.7.2	Perches must be provided by 4 weeks of age.					•	•
4.8 Environmental Enrichments							
ⓘ An enrichment is an addition to the chicken's environment that encourages the expression of natural behavior such as foraging, without losing their novelty. Examples include, but are not limited to, bales of straw or hay, raised platforms and perches, provision of forages or brassicas, and scattered grains. Dust-baths and litter are not considered enrichments. Items that quickly lose their novelty and/or go unused by the chickens are not considered acceptable enrichments. ⓘ Examples of enrichments that qualify for this section (and those that do not) are shown in Appendix VIII. This Appendix outlines that one enrichment is defined as either one item from Section A or two (2) items from Section B. ⓘ For Step 4 chickens, the standards in this Section apply if chickens are removed from pasture when climatic conditions pose a welfare risk to the chickens. ⓘ As chickens raised in Step 5 and Step 5+ systems may only be housed at night and during extreme weather conditions, indoor enrichments are not applicable.							
4.8.1	Indoor enrichments must be provided by the time the chicks are 10 days old.	•	•	•	•		
4.8.2	Indoor enrichments must be maintained throughout the life of the chickens.	•	•	•	•		
4.8.3	The indoor environment must contain at least 1 type of enrichment (See Appendix VIII) that is used by the chickens.	•					
4.8.4	The indoor environment must contain at least 2 different types of enrichments (See Appendix VIII) that are used by the chickens.		•	•	•		
4.8.5 ⓘ	For every 1,000 ft ² (93 m ²), or part 1000 ft ² (93 m ²) of indoor space there must be a minimum of 1 enrichment. ⓘ [4.8.5]: For example, for a house that is 1200 ft ² two enrichments must be provided, for a house that is 1400 ft ² two enrichments would be required and for a house that is 2020 ft ² three enrichments would be required.	•					

STANDARD		Step level					
		1	2	3	4	5	5+
4.8 Environmental Enrichments <i>Continued</i>							
4.8.6	For every 750 ft ² (70 m ²), or part 750 ft ² (70 m ²) of indoor space, there must be a minimum of 1 enrichment		•	•			
	ⓘ [4.8.6]: For example, for a house that is 1500 ft ² two enrichments must be provided, for a house that is 1800 ft ² three enrichments would be required and for a house that is 3000 ft ² four enrichments would be required.						
4.8.7	For every 500 ft ² (46 m ²), or part 500 ft ² (46 m ²) of indoor space, there must be a minimum of 1 enrichment.				•		
	ⓘ [4.8.7]: For example, for a house that is 1000 ft ² two enrichments must be provided, for a house that is 1600 ft ² four enrichments would be required and for a house that is 2200 ft ² five enrichments would be required.						
4.8.8	For houses that are smaller than 1,000 ft ² (93 m ²), there must be a minimum of 2 enrichments.	•					
4.8.9	Indoor enrichments must be distributed throughout the house.	•	•	•	•		

5 OUTDOOR CONDITIONS

- ① Outdoor access is defined as an outdoor area such as a dry lot, concrete pad, or pasture, but does not have to be pasture.
- ① Pasture includes access to rangeland, grassland, planted pastures, managed pastures, wooded areas, and any other areas where chickens have access to vegetation.
- ① Pastures should be rested and allowed to regenerate between groups.

STANDARD		Step Level					
		1	2	3	4	5	5+
5.1 Outdoor and/or Pasture Access							
① Operations are strongly encouraged to provide chickens access to the outdoors from the youngest age possible and for as much of their lives as possible.							
5.1.1	All chickens from 4 weeks of age must be given continuous outdoor access during daylight hours unless climatic conditions pose a welfare risk. ① [5.1.1]: Chickens in Step 3 systems have seasonal outdoor access.			•			
5.1.2	All chickens from 4 weeks of age must be given continuous access to pasture during daylight hours. If climatic conditions pose a welfare risk then chickens can be removed from pasture but must be given continuous access to an outdoor area during daylight hours. If the risk from climatic conditions is considered extreme, access to outdoor areas may be restricted as long as the restriction does not exceed 25 days (cumulative total) throughout a calendar year. ① [5.1.2 a]: Chickens in Step 4 systems may only be confined to housing during extreme weather conditions (e.g. heavy precipitation, tornadoes, hurricanes, monsoons, blizzards, floods or non-typical weather for the season such as large swings in temperature) that jeopardize their welfare. ① [5.1.2 b]: A calendar year is defined as January 1 st through December 31 st . ① [5.1.2 c]: See Section 5.2 for requirements for pasture and outdoor areas.				•		
5.1.3	All chickens from 4 weeks of age must be given continuous access to the pasture during daylight hours. If the risk from climatic conditions is considered extreme, access to outdoor areas may be restricted as long as the restriction does not exceed 5 consecutive days and 25 days (cumulative total) throughout a calendar year. Seasonal housing due to inclement climatic conditions is prohibited. ① [5.1.3 a]: Chickens in Step 5 and Step 5+ systems may only be confined to housing during extreme weather conditions (e.g. heavy precipitation, tornadoes, hurricanes, monsoons, blizzards, floods or non-typical weather for the season such as large swings in temperature) that jeopardize their welfare. ① [5.1.3 b]: A calendar year is defined as January 1 st through December 31 st .					•	•
5.1.4	The following records are required: a. date chickens are first given outdoor or pasture access; b. any day that outdoor or pasture access is denied; and c. reasons for any denial of outdoor or pasture access.			•	•	•	•
5.1.5	Any chickens slaughtered before 28 days of age must have access to the outdoors during daylight hours for a minimum of 2 weeks, otherwise the maximum Step-level that is possible for those flocks is a Step 2. ① [5.1.5]: See Standard 5.1.4 for record requirements.			•	•	•	•

STANDARD		Step Level					
		1	2	3	4	5	5+
5.1 Outdoor and/or Pasture Access <i>continued</i>							
5.1.6	Openings from the house, whether stationary or mobile, must allow chickens free access to the outdoors during daylight hours.			•	•	•	•
	<p>📌 [5.1.6]: As houses are constructed in a variety of designs and dimensions, this standard does not designate a minimum number of access points or the dimensions of each opening. Rather, it requires that there are sufficient entry / exit points so that chickens are readily able to access the outdoors. A house may have several smaller openings spaced along one or more walls (e.g., pop holes) or a single large opening (e.g., a door or wall segment).</p>						
5.2 Vegetation and Foraging Materials							
<p>📌 There is no set square footage for pasture space per bird (see Section 5.3 for space requirements for outdoor areas), but operations must provide sufficient space such that the vegetative cover conditions in Section 5.2 can be met throughout the life of the flock. The actual area per chicken that is required to meet the requirements of Section 5.2 throughout the life of the flock will depend on the region and/or climate where the chickens are kept.</p>							
5.2.1	At least 25% of each occupied outdoor area must be covered with vegetation and/or forage.			•			
	<p>📌 [5.2.1]: This standard can be met by placing cut or harvested vegetation and/or forages, such as alfalfa hay, in the outdoor area). Litter cannot be used as a foraging material for this standard.</p>						
5.2.2	Within 200ft (61m) from the house, at least 50% of each occupied pasture area must be covered with vegetation and/or forage accessible at chicken height throughout the life of the flock.				•		
	<p>📌 [5.2.2]: See Appendix IX for a definition and illustration of “at chicken height”.</p>						
5.2.3	Within 200ft (61m) from the house, at least 75% of each occupied pasture area must be covered with vegetation and/or forage accessible at chicken height throughout the life of the flock					•	•
	<p>📌 [5.2.3]: See Appendix IX for a definition and illustration of “at chicken height”.</p>						
5.2.4	No more than half of the vegetation required by Standard 5.2.2 (25% of the total pasture area) can be cut or harvested vegetation and/or forages such as alfalfa hay.				•		
	<p>📌 [5.2.4]: Litter cannot be used as a foraging material for this standard.</p>						
5.2.5	The requirements of Standard 5.2.3 must be met by growing vegetation and cannot be met by placing cut or harvested vegetation and/or forages in the pasture area.					•	•
5.2.6	Denuded areas cannot extend more than 10ft (3m) from the house.					•	•
5.2.7	If stationary housing is used, there must be at least two pasture areas accessible to the chickens, such that one can be rested while the other is in use.					•	•
5.2.8	If chickens are removed from pasture during winter months, the outdoor area must include materials that encourage foraging behavior.				•		
	<p>📌 [5.2.8]: Materials that encourage foraging behavior include, but are not limited to, whole grains, hay and straw.</p>						
5.3 Space Requirements for Outdoor Areas							
5.3.1	The outdoor area must be equal to or greater than 75% of the total indoor floor space of the house.			•			
5.3.2	When removed from pasture during winter months, the outdoor area must be equal to or greater than 100% of the total indoor floor space of the house.				•		

STANDARD		Step level					
		1	2	3	4	5	5+
5.4 Provisions to Encourage Chickens to Use Outdoor and Pasture Areas ⓘ If chickens feel safe, they are more likely to go outside and use the pasture or outdoor area. Provisions to achieve this are listed in Appendix X. ⓘ The house, whether stationary or mobile, does not qualify as meeting the requirements of this Section.							
5.4.1 ⓘ	Each occupied outdoor and/or pasture area must contain features that encourage the chickens to range.			•	•	•	•
	ⓘ [5.4.1]: Examples include, but are not limited to, bushes, shrubs, shade cloth, A-frame structures, and perches. The house, whether stationary or mobile, including the roof and typical 12" overhang, does not qualify as meeting the requirement of this standard. Any structures attached to the house, such as verandas or awnings, can be counted towards a percentage of the requirements in 5.4.3 as long as they provide free access to the outdoors, and are not the only provision provided (see 5.4.3 for more details). The provisions should be dispersed throughout the outdoor area to encourage roaming and foraging.						
5.4.2	Provisions not attached to the house, must be distributed in a way that encourages chickens to range, and must start within 15 ft (4.5 m) of the house.			•	•	•	•
5.4.3	For birds placed from 1 January 2018 to 31 December 2023, provisions that encourage chickens to range must provide a cumulative total of at least 8 ft ² (0.75 m ²) of cover for every 250 chickens in the flock.			•	•		
	For birds placed from 1 January 2024, onwards provisions that encourage chickens to range must provide a cumulative total of at least 8 ft ² (0.75 m ²) of cover for every 100 chickens in the flock.			•	•		
	ⓘ [5.4.3 a]: G.A.P. recognizes that operations need time to provide provisions in outdoor areas and pasture, particularly if an increase in natural shelter (shrubs, bushes, trees) is intended to meet the 2024 requirements. Operations should use the time allotted to make progress towards meeting the requirements of the 2024 standard – i.e. planting trees/shrubs now so they will provide the required cover by 2024. ⓘ [5.4.3 b]: The requirements of this Standard apply to the total number of birds in the flock, not the number of birds that are using the outdoor or pasture area at any one time. ⓘ [5.4.3 c]: Verandas or awnings attached to houses/barns can count towards a maximum of 50% of the total square footage required (e.g. a flock of 10,000 chickens requires 320 ft ² of outdoor provisions - 160 ft ² of the 320 ft ² can be a veranda attached to the house. The remaining balance of the square footage required (160 ft ²) must be natural or other artificial items not attached to the house.						
5.4.4	Provisions that encourage chickens to range must provide a cumulative total of at least 8 ft ² (0.75 m ²) of cover for every 100 chickens.					•	•
	ⓘ [5.4.4 a]: The requirements of this Standard apply to the total number of birds in the flock, not the number of birds that are using the outdoor or pasture area at any one time. ⓘ [5.4.4 b]: Verandas or awnings attached to houses/barns can count towards a maximum of 50% of the total square footage required (e.g. a flock of 2,000 chickens requires 64 ft ² of outdoor provisions - 32 ft ² of the 64 ft ² can be a veranda attached to the house. The remaining balance of the square footage required (32 ft ²) must be natural or other artificial items not attached to the house.						

6 LOADING

① *Transportation companies that are currently Poultry Handling and Transportation Certified are automatically in compliance with Standards 6.3.1 to 6.3.4, 6.3.8, 6.3.9 and 6.4.1.*

STANDARD	Step Level					
	1	2	3	4	5	5+
6.1 Condition of Chickens						
6.1.1 	Transporting unhealthy, non-ambulatory, or injured chickens is prohibited.					
6.1.2	All unhealthy, non-ambulatory, injured, or small chickens (runts) who are not loaded for slaughter must be euthanized the same day that chickens from the same flock are transported or slaughtered on-site.					
	① [6.1.2]: See Section 1.5 for euthanasia requirements.					
6.2 Water and Feed Withdrawal						
6.2.1	All chickens must have continuous access to drinking water:					
	a. until loading begins if water lines do not need to be elevated prior to catching and loading; or					
	b. until 1 hour before loading begins if water lines must be elevated prior to catching and loading.					
6.2.2	Feed must not be withheld for more than 12 hours prior to slaughter.					
6.3 Catching and Loading						
6.3.1	Lights must be dimmed throughout the catching and loading process.					
6.3.2	Chickens must be caught calmly to minimize stress and risk of injury.					
6.3.3 	Kicking, throwing, striking, punching, hitting, or otherwise causing injury to chickens is prohibited.					
6.3.4 	Chickens must never be lifted or carried by the head, neck, one or both wings, or tail.					
	① [6.3.4]: G.A.P. discourages the carrying of chickens by a single leg and urges industry to move away from this practice.					
6.3.5	Catchers are prohibited from carrying more than 4 chickens per hand.					
6.3.6	Catchers are prohibited from:					
	a. carrying more than 2 chickens per hand; and/or					
	b. carrying chickens by a single leg.					
6.3.7	Each chicken must be caught by the body with both hands and carried upright.					
6.3.8	Mechanical loaders and conveyor belts are permitted for catching and loading chickens into containers only if they are:					
	a. well-maintained;					
	b. well-managed; and					
	c. do not cause harm to the chickens.					
	① [6.3.8]: If operations at Step 5+ are using mechanical loaders this standard supersedes Standard 6.3.7.					
6.3.9	Chickens must be loaded into transport containers without causing injury.					

STANDARD		Step Level					
		1	2	3	4	5	5+
6.4 Transport Containers for Chickens							
6.4.1	Containers, whether modules, coops, drawers, or other, must be in clean and sound operational condition, and of a design that does not cause injury to the chickens.	●	●	●	●	●	●
	📌 [6.4.1]: <i>A container does not necessarily have to be washed in order for it to be clean.</i>						
6.4.2	All chickens must be able to sit on the floor of the container at the same time.	●	●	●	●	●	●
6.4.3	Containers must be of adequate height to allow the chickens to move their heads freely while sitting.	●	●	●	●	●	●

7 TRANSPORT

① The standards in this Section apply to any transport of chickens onto, within, or off of the operation. Standards for transport containers can be found in the LOADING Section.

STANDARD		Step Level					
		1	2	3	4	5	5+
7.1 Transport of Chicks							
① See also Section 2.3.							
7.1.1	A record of the total number of chicks received from the hatchery for each flock is required.	•	•	•	•	•	•
7.1.2	A record of the total number of dead-on-arrivals (DOAs) for each flock is required.	•	•	•	•	•	•
7.1.3	Mortality during transport must not exceed 0.5% per shipment.	•	•	•	•	•	•
7.1.4	Ⓜ The thermal comfort of chicks must be maintained at all times through management and the provision of supplemental heating and/or cooling, as necessary.	•	•	•	•	•	•
7.1.5	Each chick must be provided a minimum of 4 in ² (25.8 cm ²) floor space within the delivery box.	•	•	•	•	•	•
7.1.6	Each delivery box must be designed with holes for ventilation.	•	•	•	•	•	•
7.1.7	Each delivery box must contain a new, absorbent floor pad or be constructed in such a way that chicks stay dry and do not slip.	•	•	•	•	•	•
7.1.8	Chicks delivered via the mail or courier service must be in boxes prominently labelled “LIVE CHICKENS.”	•	•	•	•	•	•
7.1.9	Chicks must be delivered to the operation within 48 hours of removal from the hatcher. ① [7.1.9]: The purpose of this standard is to ensure chicks get feed and water before the resources from their yolk sac run out. Hatcheries that use the Hatch Tech system and provide food and water immediately on hatching do not have to meet this requirement.	•	•	•	•	•	•
7.1.10	Documentation of each shipment of chicks from the hatchery is required that includes: a. time chicks were removed from the hatcher; b. time of arrival onto the operation applying for certification to this standard; and c. time of placement.	•	•	•	•	•	•
7.2 Equipment and Vehicles for Chickens							
7.2.1	Equipment (e.g., a trailer) and vehicles must be managed to provide for the thermal comfort of chickens at all times.	•	•	•	•	•	•
7.2.2	If equipment (e.g., a trailer) or vehicles have open sides or tops, they must have cover(s) that can be fitted securely and adjusted as necessary to protect chickens from inclement weather.	•	•	•	•	•	•
7.3 Transport Personnel Responsibilities and Procedures							
7.3.1	Ⓜ A clear, written procedure must be made available to the auditor and/or certification company that includes actions and contact information for the driver to follow in case of accident or emergency.	•	•	•	•	•	•
7.3.2	The driver must be knowledgeable in all of his or her responsibilities and transport protocols, including those in the case of accident or emergency as per Standard 7.3.1.	•	•	•	•	•	•

STANDARD		Step Level					
		1	2	3	4	5	5+
7.3 Transport Personnel Responsibilities and Procedures <i>continued</i>							
7.3.3	If the vehicle is transporting chickens from more than one operation and/or source, each different group of chickens must be segregated in identifiable containers.	•	•	•	•	•	•
7.3.4	If chickens are transported or moved with other species on the same vehicle, they must be placed in separate compartments.	•	•	•	•	•	•
7.4 Transport Duration for Chickens							
<p>① <i>Duration of any transport is calculated for each vehicle or trailer and begins when the first chicken is loaded into the container and ends when the vehicle or trailer arrives at its destination under normal/typical driving conditions for that region. In the review of transport duration records, the certifier will take into account cases of unexpected inclement weather, vehicle accidents or malfunction, or other unforeseen circumstances that result in a delay. See Section 7.5 for records requirements.</i></p>							
7.4.1	Ⓜ Transport duration must not exceed 6 hours	•	•	•	•	•	
7.4.2	Chickens must remain on the operation at all times through slaughter, whether slaughtered in an on-site facility or a mobile unit.						•
	① [7.4.2 a]: Chickens raised in Step 5+ systems can only be moved to on-site or mobile slaughter.						
	① [7.4.2 b]: Step 5+ chickens can be crated for movement to the on-site or mobile slaughter facility. ① [7.4.2 c]: Chickens can only be transported or moved off the operation in cases of veterinary emergency or extreme weather conditions posing immediate risk to their survival. Transport or movement off the operation for either of these two reasons will not result in non-compliance with the standard.						
7.5 Transport Records							
7.5.1	The following records are required for each vehicle transporting chickens: a. date of transport; b. starting time for catching and loading chickens into each truck; c. arrival time at slaughter facility; d. reasons for any stops or delays en route; and e. number of chickens transported from the operation.	•	•	•	•	•	
7.5.2	Separate transport documentation, whether a bill of lading, delivery note, or other, is required for each operation if the vehicle is scheduled to pick up chickens from more than one operation.	•	•	•	•	•	
7.5.3	The following records are required for all chickens slaughtered on site: a. date of transport; b. starting and ending times for catching and loading chickens; c. arrival time at the place where chickens will be slaughtered; and d. number of chickens moved and slaughtered on-site.						•

8 RODENT, WILD BIRD, AND PREDATOR CONTROL

STANDARD		Step Level					
		1	2	3	4	5	5+
8.1 Rodent Control							
① <i>The standards in this Section are applicable to any rodent control efforts, whether contracted or not.</i>							
8.1.1	Good sanitation must be the first level of rodent control.	●	●	●	●	●	●
	① [8.1.1]: <i>Good sanitation includes exclusion of rodents from buildings, bays, or bins where chickens live and where grain or other feeds are stored; clear up of spills of feed; and management of trash to reduce attracting or harboring rodents.</i>						
8.1.2	If good sanitation is ineffective, an integrated rodent control program must be implemented. This program must include: <ul style="list-style-type: none"> a. methods of control that only target rodents; b. an assessment of different methods of lethal control that are commercially available; c. if traps are used they must be species specific, appropriately located, and must be designed to cause rapid death; and/or d. licensed rodenticides are only used in areas where traps will be ineffective (traps are most effective in enclosed spaces and rodent pathways) OR when monitoring is required, for example as part of salmonella reduction protocols. 	●	●	●	●	●	●
	① [8.1.2]: <i>Glue boards, drowning, and drowning traps do not meet the above requirements.</i>						
8.1.3	Multiple catch traps (e.g. tin cats) used for monitoring rodent populations must be baited with rodenticide.	●	●	●	●		
8.2 Wild Bird Control							
8.2.1	Wild birds must be excluded from housing.	●	●	●			
	① [8.2.1]: <i>This Standard only applies to Step 3 operations when chickens are excluded from outdoor areas as described by Standard 5.1.1.</i>						
8.3 Predator Control							
① <i>The standards in this Section are applicable to any predator control efforts, whether contracted or not.</i>							
① <i>Control of predators must not violate any local, state, provincial, territorial, federal, national, or other laws.</i>							
① <i>Note that certain mammals and avian species of the orders Falconiformes, Accipitriformes, Strigiformes and Cathartiformes are strictly protected and any exceptions require approval by relevant wildlife permitting authorities.</i>							
8.3.1	When predators are considered to be a problem, each operation must have a predator control program in place.	●	●	●	●	●	●
8.3.2	Non-lethal methods of predator exclusion from housing and occupied outdoor and pasture areas must be the first level of control.	●	●	●	●	●	●
8.3.3	If non-lethal methods are ineffective and chickens are at risk, shooting is the only method of lethal control allowed, and is only allowed if the shooter is skilled, and the shot kills immediately.	●	●	●	●	●	●
8.3.4	Ⓜ Poisons, drowning, all snares, leg hold traps and all traps other than live traps are prohibited.	●	●	●	●	●	●
8.3.5	Any live traps must be checked at least once daily and acted upon within 24 hours.	●	●	●	●		
	① [8.3.5]: <i>Live traps, also known as humane traps, cannot contain poison or maim, or in any other way result in lethal control.</i>						

STANDARD		Step Level					
		1	2	3	4	5	5+
8.3 Predator Control <i>Continued</i>							
8.3.6	Any live traps must be checked at least twice daily and acted upon within 24 hours.					•	•
	📌 [8.3.6]: <i>Live traps, also known as humane traps, cannot contain poison or maim, in any other way result in lethal control.</i>						
8.4 Guardian Animals							
8.4.1	Guardian animals, if used, must be:						
	<ul style="list-style-type: none"> a. well trained; b. capable of deterring predators in the area; c. does not harm the flock it is guarding; and is d. suitable for the environmental conditions of the operation. 	•	•	•	•	•	•
📌 [8.4.1]: <i>This includes dogs, llamas, donkeys and any other animals that may be used for guardian duties.</i>							

9 PLANS, PROTOCOLS, PROCEDURES, TRAINING, RECORDS AND DOCUMENTS

① Certain historical records and documents included in this section may not be available at the time of initial audit as the operation applying for G.A.P. certification was unaware they would be required and, therefore, cannot create them for past events, treatments, assessments, or practices. At the time of initial audit, record-keeping and documentation mechanisms must be place to meet each of these standards, and records and documentation for, at a minimum, the chickens presently on-site must be available.

STANDARD		Step Level					
		1	2	3	4	5	5+
9.1 General Records Requirements							
9.1.1	Records must be written and made available to the inspector and/or certification company. Acceptable formats include but are not limited to; record sheets and cards, calendars, notebooks, and computer documents.	•	•	•	•	•	•
	① [9.1.1]: Records can be collected and stored by producer groups, but must be available at the time of the audit.						
9.1.2	Records must be presented in an organized manner.	•	•	•	•	•	•
9.1.3	All records, reports, Step certificates, and other materials and correspondence relating to Step certificates must be kept for at least one certification cycle.	•	•	•	•	•	•
	① [9.1.3]: These materials must be kept until after the operation has been re-certified.						
9.2 Written Farm/Animal Health Plan/System Plan							
9.2.1	Each operation must have a written plan describing: <ul style="list-style-type: none"> a. an overview of the operation, including size, type/stage of production, location, and typical climatic conditions; b. emergency procedures, including those for natural disasters, fire, water shut off, and, if applicable, power failure; c. operational practices and policies for chicken production: <ul style="list-style-type: none"> i. provision for daily feed and water, including ration details; ii. health programs (e.g., supplementation, vaccination and other preventative, maintenance and/or health-promoting practices); iii. routine husbandry procedures; iv. care of sick and/or injured chickens, including on-farm euthanasia policies; v. management of outdoor areas, if applicable to production system; vi. brooding; and vii. rodent, wild bird, and predator control practices; and d. environmental management (i.e. to reflect how various environmental challenges are handled such as large fluctuations in temperature, excessive humidity, etc.). 						
	① [9.2.1]: The Farm and Animal Health Plan can be provided by an affiliated group (e.g., a producer group, co-operative, marketing entity) or created with the aid of external consultation (e.g., extension agents, veterinarians, peers), but must include information specific and relevant to the operation applying for G.A.P. certification.						

STANDARD		Step Level					
		1	2	3	4	5	5+
9.3 Biosecurity Procedures and Protocols							
9.3.1	<p>Each operation must have a documented and implemented biosecurity program that covers:</p> <ul style="list-style-type: none"> a. procedures for bringing any chickens onto the site, including new chicks and any returning chickens (e.g. show chickens); b. procedures and policies for caretakers (e.g. dress code and footwear requirements, policies concerning pets and/or the keeping of chickens or other chickens off-site); c. procedures and policies for visitors to the operation (e.g. minimizing visitors, provision of foot baths, booties, and protective clothing; visitor logs); d. feed trucks and equipment delivery to the operation; e. shared borders with neighboring operations, if applicable; and f. clean-out procedures of housing units between flocks. 						
	<p>i [9.3.1]: <i>G.A.P. has specific protocols in the event of an avian influenza or similar disease outbreak. Please check with your certifier in the event a localized outbreak impacts your operation's biosecurity procedures.</i></p>						
9.4 Alternative Power Supply							
9.4.1	<p>If power is essential to the operation of heating, cooling, ventilation, watering, and/or feeding systems, each operation must have:</p> <ul style="list-style-type: none"> a. an alternative power supply; and/or fail-safe device in working condition; and b. a method of notification in the event of power failure. 						
9.5 Training							
9.5.1	<p>Each operation must provide training to all care-givers (whether full-time, part-time, seasonal or contractual) and/or managers that:</p> <ul style="list-style-type: none"> a. is written and/or hands-on; b. is presented in all necessary languages; c. includes instruction on recognizing signs of normal and abnormal chicken appearance and behavior; d. describes all aspects of the individual's responsibilities; e. describes emergency procedures; f. is provided prior to the individual's handling of any chickens on the operation; g. covers all requirements of this version of G.A.P.'s 5-Step® Animal Welfare Standards for Chickens Raised for Meat; and h. is on-going as necessary and, at a minimum, when any changes affecting the care and management of chickens are implemented. 						
	<p>Each operation must keep a record of care-giver's training (whether full-time, part-time, seasonal or contractual), including dates of training and topics covered.</p> <p>i [9.5.2 a]: <i>Training includes initial, re-training and on-going training.</i></p> <p>i [9.5.2 b]: <i>This record-keeping standard applies to all care-givers but does not include immediate family members.</i></p>						

STANDARD		Step level						
		1	2	3	4	5	5+	
9.6 Traceability and Chain of Custody								
9.6.1	Each operation must have individual flock records that can trace the flock from the hatchery/hatching through to slaughter.	•	•	•	•	•	•	
	<p>① [9.6.1 a]: See 5(d) in the Program Requirements for segregation protocol requirements.</p> <p>① [9.6.1 b]: For operations supplying producer groups, it is acceptable for the producer group to record and maintain this information for individual operations.</p> <p>① [9.6.1 c]: See 1(j) in the Program Requirements for Chain of Custody Records Reconciliation Program (CCRRP).</p>							
9.6.2	<p>Each G.A.P. Certified operation must send a chain of custody record with each truck of chickens transported off the operation to the slaughter facility. The record must include:</p> <ul style="list-style-type: none"> a. the number of chickens transported; b. date of transport; c. Step-level; d. certificate number; and e. certificate expiry date. 	•	•	•	•	•		
9.6.3	A copy of the chain of custody record from each load of chickens must be kept by the operation for review by the auditor at re-inspection.	•	•	•	•	•	•	

10 SLAUGHTER REQUIREMENTS

STANDARD		Step Level					
		1	2	3	4	5	5+
<p>10.1 General Requirements</p> <p>① Operations, or the affiliated group (e.g. a producer group, co-operative, marketing entity), must provide a complete copy of the current third-party animal welfare audit for all slaughter facilities used by the Operation or affiliated group to the certifier to verify this Section. G.A.P. certifiers will review the core criteria of the current audit and assess eligibility to meet this standard (i.e. a desk audit of a current third-party animal welfare audit).</p> <p>① The animal welfare slaughter audit does not have to be conducted by a G.A.P. approved certification company. Operations and suppliers are encouraged to check with their retail partners to ensure the certifier used for their slaughter audit is in compliance with the retailer’s buying requirements.</p> <p>① A Robust Systematic Humane Handling and Slaughter Plan is not equivalent to a third-party animal welfare audit and does not replace the need for such a third-party audit.</p>							
10.1.1	Birds must be stunned and rendered insensible prior to slaughter.	•	•	•	•	•	•
10.1.2	<p>All operations holding a current G.A.P. certificate, where birds will be marketed as G.A.P. Certified, must be slaughtered at a slaughter facility that has passed and holds a current annual third-party animal welfare audit with a minimum score of 90% and pass all core criteria listed below:</p> <ul style="list-style-type: none"> a. at least 99% of the chickens must be properly stunned (on a sample of 500); b. no more than 1% broken or dislocated wings; c. if hung live, both legs must be shackled (1 in 500 is a failure); d. no broken legs (1 broken leg in 500 is a failure); e. no more than 1% bruised thighs and drumsticks; and f. no live chickens entering the scalders. <p>Source: adapted from the Slaughter section of National Chicken Council Animal Welfare Guidelines and Audit Checklist for Broilers (2014).</p> <p>① [10.1.2]: Any bird found with a broken leg or wing should be euthanized immediately, rather than shackled.</p>	•	•	•	•	•	•
10.1.3	Chickens that are dead-on-arrival (DOA) must not exceed 0.5%.	•	•	•	•	•	•
10.1.4	<p>In addition to the criteria listed in 10.1.1 – 10.1.3, third-party slaughter facility audits must include, at a minimum, the following information as part of the audit form(s):</p> <ul style="list-style-type: none"> a. auditor name; b. audit date; c. full name, address, and contact information of slaughter facility; d. establishment number; e. name of slaughter facility representative who accompanied the auditor on the audit; f. summary table of audit findings by section (i.e. lists and scores for core criteria such as effective stunning, insensibility on the bleed rail, vocalization etc. as well as secondary criteria); g. type of slaughter system (e.g. electrical water bath, CAS etc.); and h. typical line speed of slaughter facility (if applicable). 	•	•	•	•	•	•

STANDARD		Step level					
		1	2	3	4	5	5+
10.2 Segregation and In-Plant Traceability							
10.2.1	All slaughter facilities used by the operation or affiliated group must provide a copy of their written in-plant segregation and traceability protocol outlining how G.A.P. Certified chickens, and differing Step-levels if applicable, are kept segregated from non-G.A.P. Certified chickens and product.	•	•	•	•	•	•
10.2.2	Chickens must be slaughtered on-farm using an on-farm slaughter facility or a mobile slaughter unit.						•

Appendix I: Breeds/Strains Growth Rates and Step-levels

Section 1.2 sets standards on the maximum growth rate that is acceptable for each Step level. G.A.P. has used published breeders' data to assess the average daily gain (as hatched) of a variety of commonly used breeds/strains and therefore the maximum Step-level that can be achieved when using that breed/strain. This information is shown in the table below.

The maximum Step-level achievable by each breed is set from the standardized data, reviewed by G.A.P., for chickens at 42 days of age. G.A.P. understands that some operations achieve different growth rates to those shown below and that some operations rear birds to different slaughter ages. However, this is not relevant for the purposes of this standard. As noted at the start of Section 1 of these standards, G.A.P. announced in March 2016 that it would be transitioning all Step Levels to higher welfare genetics by January 1, 2024. To assist us to that end, G.A.P. has commissioned a study to determine the specific requirements for each Step level. Until that study is complete, G.A.P. has set a standard way of assessing breeds and strains and the Step level for which each is suitable. The table below is the result of this process. In practice this means, for example, a Step 3 operation can use a Cobb 500 bird regardless of their on-farm performance and growth rate and their age/weight at slaughter.

If an operation wishes to use an alternative breed/strain they should contact their Certifier for confirmation that the bird they wish to use meets the requirements of their expected Step-level.

Source	Breed/Strain	Av Daily Gain (as hatched) (lbs /Day)	Maximum Step Achievable
Cobb Vantress	Cobb 500	0.150	3
Cobb Vantress	Cobb Sasso 150	0.077	5+
Cobb Vantress	Cobb 700	0.139	3
Ross	Ross 308	0.147	3
Ross	Ross 708	0.140	3
Ross	Rowan Ranger	0.090	5
Hubbard	Classic	0.150	3
Hubbard	F15	0.147	3
Hubbard	Flex	0.150	3
Hubbard	H1	0.149	3
Hubbard	JV	0.140	3
Hubbard	F915	0.136	3
Hubbard	JA287	0.110	4
Hubbard	JA787	0.110	4
Hubbard	JA987	0.110	4
Hubbard	JA257	0.099	5

....table continues on next page

Source	Breed/Strain	Av Daily Gain (as hatched) (lbs/day)	Maximum Step Achievable
Hubbard	JA757	0.099	5
Hubbard	JA957	0.099	5
Hubbard	JACY87	0.099	5
Hubbard	JACY57	0.088	5
Hubbard	Redbro	0.077	5+
Hubbard	Redbro Cou Nu (Poulet Rouge Fermier)	0.065	5+
Crystal Lake Farms	Free Ranger	0.091	5
Freedom Ranger Hatchery	Freedom Ranger	0.092	5
Centurion Poultry	Sagitta	0.048	5+
S&G Poultry	Heritage White	0.095	5
S&G Poultry	Red & Rainbow Ranger	0.074	5+

Appendix II: Identifying and Scoring Footpad Dermatitis

Section 2.8 requires a footpad dermatitis monitoring program, conducted at the slaughter facility or on-farm in the last week prior to slaughter, that includes random sampling. One foot is assessed from each selected chicken. The total number of chickens selected depends on the number of flocks being sent to slaughter on a particular day – see ‘Determining the Sample Group’ below. The identification and scoring protocol below considers both the size and severity of lesions. If 100 birds are assessed the total maximum footpad dermatitis score would be 200 (i.e., all 100 feet scoring ‘2’). The maximum thresholds for a flock’s total footpad dermatitis sum are listed in standards 2.8.3 to 2.8.6 according to Step level.

Standards 2.8.3 to 2.8.6 set limits for footpad dermatitis as an annual average. This means that an operation with some seasonal variation in footpad dermatitis may be within the standards on an annual basis. For example, a Step 2 operation sends one flock to slaughter every month. The scores for each flock were as follows: January = 20; February = 18; March = 15; April = 15; May = 10; June = 10; July = 10; August = 10; September = 12; October = 12; November = 14; December = 14. The annual average score is therefore 13.5. Standard 2.8.3 requires Step 2 operations to have an annual average footpad score of 15 so the operation is in compliance with this standard.

However, standard 2.8.7 requires an intervention plan (see Appendix III) to be written and implemented for each individual flock that exceeds the level set in Standard 2.8.3. This operation must therefore have written a plan to reduce their risk of footpad dermatitis in January of the example year above and created a new plan, updating their progress and detailing any new actions, in February of the example year.

PROTOCOL

Step 1: Determining the Sample Groups

- At the slaughter facility: Determine how many flocks were sent from each operation. The minimum sample size is 100 birds if only one flock is sent to slaughter. If more than one flock is sent to slaughter an additional 25 birds for each additional flock must be assessed; for example, if 5 flocks are sent to slaughter on the same day the sample group must be 100 plus 25, plus 25, plus 25, plus 25 for a total of 200 birds assessed with one foot for each bird scored. Once the sample size is set, randomly collect a total of different legs to meet the sample size, and score the footpads at a convenient time during the day.
- On-farm: Determine how many flocks will go to slaughter on the same day. The minimum sample size is 100 birds if only one flock is sent to slaughter. If more than one flock is sent to slaughter an additional 25 birds for each additional flock must be assessed; for example if 3 flocks are sent to slaughter on the same day the sample group must be 100 plus 25, plus 25 for a total of 150 birds assessed with one foot for each bird scored. Once the sample size is set, in the last week before those birds go to slaughter, randomly catch chickens in 4 different locations in each house to meet the required sample size, examine their feet and score the footpads. A representative sample of birds across all flocks must be assessed. If three flocks are sent to slaughter and 150 in total must be scored, 50 should be scored in each house.

Appendix II: Identifying and Scoring Footpad Dermatitis *continued*










Step 2: Scoring

- Use the following chart to score the feet from each flock with a footpad dermatitis score of '0,' '1,' or '2.'

Step 3: Calculating the Flock's Total Footpad Dermatitis Sum

- $(\# \text{ of feet scoring '0'} \times 0) + (\# \text{ of feet scoring '1'} \times 1) + (\# \text{ of feet scoring '2'} \times 2) / (\text{number of feet scored}/100) = \text{Total Footpad Dermatitis Sum}$
- In the above calculation, the Total Footpad Dermatitis Sum is determined as follows:
 - a foot scoring '0' does not get any points
 - a foot scoring '1' gets 1 point
 - a foot scoring '2' gets 2 points
- For example:
 - Of 100 feet from 100 chickens assessed: 65 score '0,' 30 score '1,' and the remaining 5 score '2.'
 - $(65 \times 0) + (30 \times 1) + (5 \times 2) / (100/100) = 0 + 30 + 10 / 1$, for a Total Footpad Dermatitis Sum of 40
 - Of 500 feet from 500 chickens assessed 320 score '0', 140 score '1' and the remaining 40 score '2'
 - $(320 \times 0) + (140 \times 1) + (40 \times 2) / (500/100) = 0 + 140 + 80 / 5$ for a Total Footpad Dermatitis Sum of 44

Appendix II: Identifying and Scoring Footpad Dermatitis *continued*

Score: 0	Score: 1	Score: 2
(0 points each for score of 0)	(1 point each for a score of 1)	(2 points each for a score of 2)
<ul style="list-style-type: none"> • No lesions or very small • No discoloration or slight on a limited area • No hyperkeratosis or mild • Old or no scars 	<ul style="list-style-type: none"> • Mild and/or superficial lesions • Substantial discoloration on the footpad • Dark papillae, no ulceration 	<ul style="list-style-type: none"> • Severe and significant lesions • Ulceration • Dark papillae and ulceration • Abscesses and/or swollen feet (bumble foot)
<p data-bbox="359 464 470 488">no lesion</p>  <p data-bbox="296 808 533 833">small discoloration</p>  <p data-bbox="275 1149 554 1174">completely healed scar</p> 	<p data-bbox="978 464 1119 488">mild lesion</p>  <p data-bbox="894 808 1203 833">substantial discoloration</p>  <p data-bbox="873 1149 1224 1174">dark papillae, no ulceration</p> 	<p data-bbox="1598 464 1759 488">severe lesion</p>  <p data-bbox="1503 808 1854 833">dark papillae and ulceration</p>  <p data-bbox="1482 1149 1875 1174">abscess, swollen (bumble foot)</p> 

Adapted and reprinted with permission: "Foot-pad dermatitis in broilers: a photo guide to broiler foot health classification." Dr Lotta Berg, Swedish University of Agricultural Sciences, Sweden
 G.A.P.'s 5-Step® Animal Welfare Standards for Chickens Raised for Meat | Issued 3 April 2018 | Updated 1 May 2020 | v3.2
 ©2020 Global Animal Partnership. All rights reserved, including the right to reproduce this publication or portions thereof in any form whatsoever.

Appendix III: Intervention Plans

Standards 2.8.7, 2.9.7, 2.10.6 and 4.4.3 all require written intervention plans to be put into effect if the levels of foot pad dermatitis detailed in Standards 2.8.3 (Step 1), 2.8.4 (Steps 2 and 3), 2.8.5 (Steps 4 and 5) and 2.8.6 (Step 5+); lameness detailed Standards 2.9.3 (Step 1), 2.9.4 (Steps 2 and 3), 2.9.5 (Steps 4 and 5) and 2.9.6 (Step 5+); flock mortality detailed in Standards 2.10.2 (Step 1), 2.10.3 (Steps 2 and 3) and 2.10.4 (Steps 4, 5 and 5+); and air quality detailed in Standard 4.4.2 (all Steps), do not meet the requirements of the standards. The intervention plan template below must be completed and implemented for each flock when this occurs. G.A.P. Certifiers will review these plans at audit.

Template:

Intervention plan questions	Operation's response
Which standard does the flock not meet and what level of foot pad dermatitis/lameness /mortality/air quality has been found?	
Why has the problem occurred?	
What actions have been put in place to resolve the issue now?	
What actions have been put in place to ensure this does not happen again with future flocks?	

Example of completed template:

Intervention plan questions	Operation's response
Which standard does the flock not meet and what level of foot pad dermatitis/lameness /mortality/air quality has been found?	<i>Standard 2.8.5 – the foot pad dermatitis score for one of our Step 4 flocks was 10</i>
Why has the problem occurred?	<i>We had a water leak in the house and litter quality was reduced with some areas of wet and capped litter over a wide area.</i>
What actions have been put in place to resolve the issue now?	<i>We repaired the water leak and checked water systems in our other houses. We added plenty of fresh clean litter to improve litter quality.</i>
What actions have been put in place to ensure this does not happen again with future flocks?	<i>We already make sure our ventilation program keeps temperature and humidity at acceptable levels to help maintain litter quality. Our feed program is checked by a nutritionist and we are sure this incidence of higher than usual FPD is not related to nutrition. We are slightly lower stocked than permitted by G.A.P. standards so we do not think stocking density is a contributory factor in this case either. We are taking more care of water line maintenance and adjusting the height of water lines as birds grow to reduce spillage and water wastage.</i>

This template is also available for download on our website on the Producer Resources page.

Appendix IV: 3-Category Lameness Evaluation

Standard 3.15 requires a lameness evaluation conducted on-farm during the final week prior to slaughter. This 3-category lameness evaluation assesses and scores the mobility of chickens observed in 4 different locations within each house, on each floor if multi-level, or in each outdoor area. If 100 birds are assessed, the total maximum lameness sum possible is 200 (i.e., all 100 chickens scoring '2'). The maximum thresholds for a flock's total lameness sum are listed in Standards 3.16.1 to 3.16.3 according to Step level.

PROTOCOL

Step 1: Determining the Sample Groups

- Determine how many flocks will go to slaughter on the same day. The minimum sample size is 100 birds if only one flock is sent to slaughter. If more than one flock is sent to slaughter an additional 25 birds for each additional flock must be assessed; for example, if 3 flocks are sent to slaughter on the same day the sample group must be 100 plus 25, plus 25 for a total of 150 birds assessed for lameness.
- Once the sample size is set, in the last week before those birds go to slaughter, assess groups of chickens in 4 different locations in each house to meet the required sample size. A representative sample of birds across all flocks must be assessed. If three flocks are sent to slaughter and 150 in total must be scored, 50 should be scored in each house.

Step 2: Scoring and Suggested Best Practices

- Use the following chart to score chickens from each flock with a lameness score of '0,' '1,' or '2.'
- Watch one chicken at a time for at least 15 seconds.
- Each chicken's mobility must be assessed and scored from the side in order to observe foot placement and length of stride.
- Gentle nudging may be necessary to encourage movement.
- Chickens must always be approached from behind, if movement must be encouraged.
- Once the chicken is moving, back away to assess the gait and assign a lameness score of '0,' '1,' or '2' according to the chart below.

Step 3: Calculating the Flock's Total Lameness Sum

- $(\# \text{ of chickens scoring '0' } \times 0) + (\# \text{ of chickens scoring '1' } \times 1) + (\# \text{ of chickens scoring '2' } \times 2) / (\text{number of assessed birds}/100) = \text{Total Lameness Sum}$
- In the above calculation, the Total Lameness Sum is determined as follows:
 - a chicken scoring '0' does not get any points
 - a chicken scoring '1' gets 1 point each
 - a chicken scoring '2' gets 2 points each

Appendix IV: 3-Category Lameness Evaluation *continued*

- For example:
 - Of 100 chickens assessed: 62 score '0,' 23 score '1,' and the remaining 15 score '2'
 - $(62 \times 0) + (23 \times 1) + (15 \times 2) / (100/100) = 0 + 23 + 30 / 1$, for a Total Lameness Sum of 53
 - Of 300 chickens assessed: 200 score '0', 88 score '1' and the remaining 12 score '2'
 - $(200 \times 0) + (88 \times 1) + (12 \times 2) / (300/100) = 0 + 88 + 24 / 3$ for a Total Lameness Sum of 37.3

Score: 0	Score: 1	Score: 2
(0 points each for score of 0)	(1 point each for a score of 1 if one or more of the descriptions below are seen)	(2 points each for a score of 2 if one or more of the descriptions below are seen)
<ul style="list-style-type: none"> • Foot may or may not curl when lifted by the chicken • Smooth gait typically with even steps that may be uneven at times • Well-balanced • Able to walk quickly and/or run • Difficult to identify any abnormality when walking or running 	<ul style="list-style-type: none"> • Uneven gait • Foot does not curl when lifted by the chicken • Irregular, short strides • Poor balance • The chicken: <ul style="list-style-type: none"> ○ may use one or both wings to help balance while walking ○ squats within 15 seconds of standing or being forced to move by gentle nudging ○ may lie down after several steps 	<p>The chicken:</p> <ul style="list-style-type: none"> • is reluctant or unable to move, or shuffles on the ground if forced to move by gentle nudging • uses wings to help with movement • takes at most a few steps, if any
Normal	Moderately Lam	Severely Lam

Appendix V: Soiled Feather Assessment

Standard 4.3.4 requires that litter within housing structures to be of quality and quantity to (1) provide a comfortable environment and (2) allow for dust-bathing behavior. At the time of audit, auditors will perform the following Soiled Feather Assessment as an additional indicator of litter conditions by observing a minimum of 100 chickens in the flock. In addition, as part of Standard 4.3.5, operations must perform this same assessment along with a Litter Quality Assessment (see Standard 4.3.6 and Appendix VI) when the flock is between 15-20 days.

If the Soiled Feather Assessment at flock age of 15-20 days shows that more than 10% of sampled birds are severely soiled; OR the Litter Quality Assessment carried out at the same time shows that litter scores 2 or 3, Standard 4.3.7 requires that a written intervention plan is put into place to improve litter quality and Standards 4.3.8 and 4.3.9 require that subsequent Soiled Feather and Litter Quality Assessments are carried out in the same flock in the week before slaughter.

Protocol for Operations:

- a. Assess each flock between 15 and 20 days.
- b. For each assessment randomly select 10 chickens from 5 different locations throughout the house or pasture area for a total of 50 chickens.
- c. Use the scoring system below and visually assess the back and top of the chicken, not including the legs and feet. IT IS NOT NECESSARY FOR BIRDS TO BE PICKED UP TO CARRY OUT THIS ASSESSMENT.
- d. Record the assessments. For your convenience, a template score sheet has been created in case it is needed¹.
- e. If any flock has an assessment where more than 10% of sampled birds are severely soiled; OR if the Litter Quality Assessment in Appendix VI gives litter scores of more than 1, each flock must additionally be assessed during the week prior to slaughter.

Scoring (see also the photos on following page)	0 =	Clean The bird is clean or has light soiling
	1 =	Significantly soiled Feathers have large areas of dirt and/or feathers are wet and dirty on up to 50% of the back/top
	2 =	Severely soiled Feathers have large areas of dirt and/or feathers are wet and dirty on over 50% of the back/top

¹ Scoring sheets illustrated below can be downloaded from www.globalanimalpartnership.org.

BACK / TOP OF CHICKEN

Score 0: Clean to Lightly Soiled

The bird is clean or has minor areas of dirtiness

Score 1: Significantly Soiled

Feathers have large areas of dirt and/or feathers are wet and dirty on up to 50% of the back/top

Score 2: Severely Soiled

Feathers have large areas of dirt and/or feathers are wet and dirty on over 50% of the back/top



Photos courtesy of 2 Sisters Food Group Ltd. Scoring guide adapted with permission from the Royal Society for the Prevention of Cruelty to Animals (Welfare Standards for Chickens, 2011)

This template is also available for download on our website on the Producer Resources page.

SOILED FEATHER ASSESSMENT

global animal partnership

Name: _____ Flock Age (in days): _____
 House ID: _____ Flock Size (in birds): _____
 Paddock ID: _____ Breed/Strain of chickens: _____

Score 0 Clean to Lightly Soiled	Score 1 Significantly Soiled	Score 2 Severely Soiled
The bird is clean or has minor areas of dirtiness.	Feathers have large areas of dirt and/or feathers are wet and dirty on up to 50% of the back/top.	Feathers have large areas of dirt and/or feathers are wet and dirty on over 50% of the back/top.

Notes:
 Assess each bird between 15 and 30 days of age.
 Randomly select 5 chickens from 10 different locations throughout the house or pasture for a total of 50 chickens.
 Visually assess the back and top of the chickens, not including the legs and feet.
IT IS NOT NECESSARY FOR BIRDS TO BE PICKED UP TO CARRY OUT THIS ASSESSMENT
 But it needs to be the corresponding Soiled Feather Score chosen.

Chicken #	Soiled Feather Score	Chicken #	Soiled Feather Score	Chicken #	Soiled Feather Score
Location 1					
1		11		21	
2		12		22	
3		13		23	
4		14		24	
5		15		25	
Location 2					
6		16		26	
7		17		27	
8		18		28	
9		19		29	
10		20		30	
Location 3					
11		21		31	
12		22		32	
13		23		33	
14		24		34	
15		25		35	
Location 4					
16		26		36	
17		27		37	
18		28		38	
19		29		39	
20		30		40	

Total number of Score 2 birds: _____

QUESTIONS:
 # of total of the following responses:
 A) 1 or more birds in a flock have a soiled feather score of 2.
 B) The soiled feather assessment gives three scores of response 1.
 Response 0 is the baseline.
 1) Complete an observational plan to improve flock quality and ensure that feather culling does not deteriorate flocks.
 2) Conduct an additional soiled feather assessment for the flock 30 days after the original.
 3) Conduct an additional Clean Quality Assessment for the flock 30 days after the original.

This template is provided by Global Animal Partnership for use as a producer's own record keeping system. It is not intended to be used for regulatory compliance purposes. For more information on the program, please contact: GlobalAnimalPartnership.com

Appendix VI: Litter Quality Assessment

Litter serves several important functions. For example, it absorbs excess moisture from drinkers and the birds' droppings; it dilutes fecal matter so reducing contact between chickens and manure – so promoting bird cleanliness; and, it provides a protective cushion between the birds and the floor. Good litter management is therefore crucial for chicken welfare.

At the time of audit, auditors will perform the following Litter Quality Assessment. In addition, as part of Standard 4.3.6, operations must perform this same assessment along with a Soiled Feather Assessment (see Standard 4.3.5 and Appendix V).

If the Soiled Feather Assessment at flock age of 15-20 days shows that more than 10% of sampled birds are severely soiled; OR the Litter Quality Assessment carried out at the same time shows that litter scores more than 1, Standard 4.3.7 requires that a written intervention plan is put into place to improve litter quality and Standards 4.3.8 and 4.3.9 require that subsequent Soiled Feather and Litter Quality Assessments are carried out in the same flock in the week before slaughter.

Protocol for Operations:

- a. Assess each flock between 15 and 20 days of age.
- b. For each assessment pick up three litter samples from three different areas in the house – one from each end of the house and one in the middle. There is no need to dig out a sample – it can be collected from the surface.
- c. Use the scoring system below to assess litter quality. The worst score of the three samples must not exceed 1.
- d. Record the assessments. For your convenience, a template score sheet has been created in case it is needed.
- e. If any flock has an assessment at 15-20 days where litter scores are more than 1; OR if the Soiled Feather Assessment in Appendix V shows that more than 10% of sampled birds are severely soiled, an intervention plan as described in Standard 4.3.7 must be implemented AND that flock must additionally be assessed during the week prior to slaughter.
- f.

Scoring Litter Quality			Action Required?
0	DRY	Dry and crumbly, free flowing on compression (will not form a ball or clod)	No action required for this standard, but note this could be causing dusty conditions
1	MOIST	Sticky on hand when compressed, clod forms but does not hold – crumbles when released from compression	No - this is the target score
2	WET	Forms a clod when compressed that holds form when released from compression	Yes – needs work to improve litter quality
3	VERY WET	Litter is noticeably wet, moisture comes out of it on compression.	

Appendix VI: Litter Quality Assessment *Continued*

Other resources:


<https://www.poultryventilation.com/sites/default/files/tips/2011/vol23n3.pdf>

<http://www.thepoultrysite.com/articles/388/litter-quality-and-broiler-performance/>

<http://extension.uga.edu/publications/detail.cfm?number=B1267>

This template is also available for download on our website on the Producer Resources page.

LITTER QUALITY ASSESSMENT



Date: _____ Flock age (in days): _____
 House ID: _____ Flock size (when placed): _____
 Flock ID: _____ Litter type: _____

Score 0	Score 1	Score 2	Score 3
Dry	Moist	Wet	Very wet
Dry and crumbly, free flowing on compression (will not form a ball or clod)	Sticky on hand when compressed, clod forms but does not hold - crumbles when released from compression	Forms a clod when compressed that holds form when released from compression	Litter is noticeably wet, moisture comes out of it on compression
No action required for this standard, but note this could be reaching dusty conditions	No action required - this is the target score	Yes - needs work to improve litter quality	

Protocol:
 Assess each flock between 15 and 20 days of age
 Pick up three litter samples from three different areas in the house - one from each end of the house and one in the middle
 There is no need to dig out a sample - it can be collected from the surface
 Use the scoring system above to assess litter quality
 Record the scores in the table below

Area	1: Top of house	2: Middle of house	3: Bottom of house
Litter Quality Score			

GUIDANCE
 If either of the following happens:
 01 The Litter Quality Assessment gives litter scores of more than 1
 02 3 or more birds in a flock have a Sublet Feather Score of 2
 Then you must do the following:
 15 Complete an Intervention plan to improve litter quality and ensure that feather pulling does not deteriorate further
 21 Conduct an additional Litter Quality Assessment for the flock in the week prior to slaughter
 22 Conduct an additional Sublet Feather Assessment for the flock in the week prior to slaughter

This template was created by Global Animal Partnership to provide a free record keeping resource for operations seeking GAP certification. If you want to use this template on other sites, please contact GAP at standard@globalanimalpartnership.com

Appendix VII: Sensory Evaluation of Air Quality

Section 4.4 requires assessment of air quality using ammonia meters, testing strips or sensory evaluation. Most operations do not use calibrated equipment to assess air quality on a daily basis. While this type of specialized equipment is an accurate way to measure air quality, it's typically too costly for everyday use. Sensory evaluation or testing strips are more commonly used. Protocols for all three methods are shown below.

Option 1: Sensory Evaluation

Below is a subjective score that can be used to assess air quality during daily monitoring.

Scores 2-5 (Moderate, Strong, Very Strong, and Overpowering) indicate that ammonia and dust are excessive and need to be addressed immediately for both human safety and animal health.

1. Once each day during flock monitoring, score and record the air quality just prior to leaving the house.
2. Record your air quality score.
3. If action is required (score 2-5), make any necessary adjustments to ventilation, management, etc. to improve air quality in your barn(s).

Scoring Air Quality			Action Required?
0	ZERO	odor and dust not noticeable (easy to breathe)	No - acceptable air quality
	1	odor and dust hardly noticeable	
2	MODERATE	odor and dust distinct, annoying (watery eyes and/or coughing)	Yes – needs work to improve air quality
3	STRONG	odor and dust irritating (stinging eyes and mouth, and/or excessive coughing)	
4	VERY STRONG	odor and dust bearable (stinging eyes and mouth, excessive coughing, and/or pain when swallowing)	
5	OVERPOWERING	odor and dust unbearable, you need to leave the barn (hurts to breathe in)	

Option 2a: Use of ammonia test strip papers

1. Once each day during flock monitoring, score air quality just prior to leaving the house.
2. Wet the test strip with distilled water (tap water contains impurities that can affect the test result). Wave the strip in the air at bird height for 15 seconds.
4. Compare the color of the exposed test strip to the calibrated color chart provided by the manufacturer to find the ppm of ammonia in the air.
5. Record the ammonia level found.
6. If action is required (score more than 20 ppm), make any necessary adjustments to ventilation, management, etc. to improve air quality in your barn(s).

Option 2b: Use of continuous ammonia sensor papers

Some types of ammonia sensor use a detection system similar to paper test strips, but rather than testing each day these are designed to provide readings for up to two months. These sensors provide a 3-hour time-weighted average of ammonia levels on a continuous basis. The visual sensor changes color (from golden yellow to blue) depending on the ammonia concentration in the surrounding environment.

1. Note expiration date (two months from the date of first use) on the sensor.
2. Attach sensor centrally & directly above the area to be monitored with string or twine. (It is recommended to have one sensor every 5,000 ft² - approximately 70ft x 70ft).
3. Once each day during flock monitoring, score air quality just prior to leaving the house.
4. Compare sensor color (center square) to surrounding color chart (outer circle) and record the ammonia level found.
5. If action is required (score more than 20 ppm), make any necessary adjustments to ventilation, management, etc. to improve air quality in your barn(s).
6. Replace after two months of use.

Option 3a: Use of electronic ammonia meters

1. Once each day during flock monitoring, turn the ammonia meter on outside the house.
3. Go inside and walk through the house for at least 2.5 minutes.
4. Read the result from the meter.
5. Record the ammonia level found.
6. If action is required (score more than 20 ppm), make any necessary adjustments to ventilation, management, etc. to improve air quality in your barn(s).

Option 3b: Use of ammonia detection tube meters

1. Once each day during flock monitoring, score air quality just prior to leaving the house.
2. Break the ends of the single use glass tube provided with the gas detection meter.
3. Place the tube inside the hand-operated pump.
4. Pull back the handle of the pump and draw a 100 mL sample of air into the glass tube from bird height. The ammonia in the air reacts to chemicals in the tube, changing the color of the chemicals the length of the tube from purple to beige.
5. The higher the level of ammonia in the house, the longer the color change down the length of the tube. Read the result from the graduated scale on the outside of the detection meter.
6. Record the ammonia level found.
7. If action is required (score more than 20 ppm), make any necessary adjustments to ventilation, management, etc. to improve air quality in your barn(s).
8. For a more accurate reading multiple samples should be taken.

Appendix VIII: Environmental Enrichments

Chickens, like other animals, benefit from a rich environment that is stimulating and allows for them to engage in natural behavior; particularly foraging behavior and exercise. Providing enrichments can improve the welfare of chickens in a commercial setting. Section 4.8 requires Step 1-4 chickens to be provided with environmental enrichments. The following document details the importance of these items, as well as outlining enrichments that count towards the requirements of the Standards as well as those that do not.

Enrichments are an addition to the chickens' environment that encourages the expression of natural behavior such as ground scratching, pecking, and foraging. As well, enrichments that increase physical activity and promote exercise can minimize undesirable and even harmful behavior, including aggression, feather-pecking, cannibalism, flightiness, and distress. (Items that are fundamental to the welfare, including health, of the chickens, such as dust-baths and litter are not considered to be enrichments as they are basic requirements). Of course, enrichments should also ensure they do not put chickens at risk of injury or stress.

The *aim* of an enrichment, however, is to:

1. add stimuli and long-term novelty to the chickens' environment;
2. evoke – and maintain – their interest, and;
3. improve their physical, behavioral, and/or mental well-being.

Enrichments can benefit chickens raised in any setting, whether exclusively indoors, with outdoor access, or on pasture. By introducing these interactive elements, the lives of chickens can most certainly be enhanced. However, not all enrichments are the same in terms of how well they actually do “enrich” the chickens environment.

EXAMPLES OF ENRICHMENTS THAT COUNT AND DO NOT COUNT TOWARDS THE REQUIREMENTS OF SECTION 4.8

The tables below list enrichments that count towards the requirements of the Section 4.8 and those that do not, and are by no means exhaustive. The examples and discussion on why, or why not they are enrichments that count towards the requirements of Section 4.8 are intended to help understand what provisions are most meaningful to chickens.

The enrichments that count towards the requirements of the Section 4.8 are categorized into two different sections – A and B depending on the size of the enrichment and the number of ways that the chickens can interact with it, so that there is equivalence between the different choices for enrichment.

For enrichments where dimensions have been listed, operations are able to utilize smaller sized enrichments as long as the cumulative total of enrichment square footage or linear space is provided to the birds.

ENRICHMENTS FOR CHICKENS THAT MEET THE REQUIREMENTS OF SECTION 4.8 – SECTION A

1 enrichment = one item from the enrichments listed in this section (See also Section 4.8 for further details on number of enrichments by size of house)

Bales
(turkey poults in top center picture)



What is the product?

Bales of straw or hay

Why is it an enrichment?

Bales of straw or hay promote physical activity, encourage pecking and foraging behavior, provide a roosting area, stimulate the birds' curiosity, and more. Chickens are able to improve their leg health by jumping on and off the bales, as well as satisfy their pecking and foraging needs, as they interact with and manipulate the bales with their beaks. These enrichments provide an interesting addition to the birds' environment and encourage them to explore and investigate.

What are the criteria for acceptable use:

Straw bales (minimum size 2.5 ft x 1.5 ft) can be placed directly on the litter and left to be pulled apart by the chickens. Leaving the bales with their original ties (2-3 strings wrapped around the bale) typically results in longer use by the chickens. In order to count as 1 enrichment, bales must remain their original size and not be untied with flakes scattered throughout the barn (i.e. a straw bale of 10 flakes = 1 enrichment, not 10).

ENRICHMENTS FOR CHICKENS THAT MEET THE REQUIREMENTS OF SECTION 4.8 – SECTION A *continued*

1 enrichment = one item from the enrichments listed in this section (See also Section 4.8 for further details on number of enrichments by size of house)

**Spreading whole grains
(laying hens pictured)**



Photo: <http://buildachickencoopeasy.com/feeding-chickens/>

What is the product:

Scattering whole grains.

Why is it an enrichment:

Chickens enjoy a wide range of foods and have a strong drive to forage, scratch, and peck. Introducing grains through a number of different ways—such as scattered loosely allows chickens to actively engage in seeking out food in a more natural way.

What are the criteria for acceptable use:

At least 1% of the chickens' total daily diet must be provided in this form and given once a day. For example, if 100lbs (45kg) of feed is given to the flock daily, at least 1lb (0.45kg) of grains must be scattered so large numbers of birds can access them each day.

Grubs and other insects



Photo: <https://www.youtube.com/watch?v=MfnyFFIrcPo>

What is the product:

Offering grubs and other insects to the chickens.

Chickens enjoy a wide range of foods and have a strong drive to forage, scratch, and peck. Grubs and insects are part of the natural diet of chickens and have the additional benefit of providing essential amino acids. Black soldier fly larvae can be raised on waste vegetables and other organic matter.

What are the criteria for acceptable use:


Whole grubs or insects must be made available to chickens at least daily for this to be an acceptable enrichment. At least 1% of the chickens' total daily diet must be provided in this form. For example, if 100lbs (45kg) of feed are given to the flock daily at least 1lb (0.45kg) of grubs must be provided.

The grubs can either be scattered or must be presented in a way that large numbers of chickens can access the grubs.

ENRICHMENTS FOR CHICKENS THAT MEET THE REQUIREMENTS OF SECTION 4.8 – SECTION A *continued*

1 enrichment = one item from the enrichments listed in this section (See also Section 4.8 for further details on number of enrichments by size of house)

Perches



Source: <https://www.cambridge.org/core/journals/animal/article/influence-of-providing-perches-and-string-on-activity-levels-fearfulness-and-leg-health-in-commercial-broiler-chickens/22316C47DA193FA25D4BFB7A3F7FCAB5#fndtn-supplementary-materials>

What is the product:

Providing perches.

Why is it an enrichment:

Perching is a natural behavior for chickens and providing perches encourages the birds to exercise and can help strengthen their legs.

What are the criteria for acceptable use:

A length equivalent to at least 2m (6.56 ft) linear perch for every 1000 birds counts as a single enrichment.

Steps 1, 2, and 3: Perches must not be sited more than 10 inches above ground level.

Steps 4, 5, and 5+: Multi-level (>10 inches) perches can be used where birds are able to use perches for the duration of their life.

ENRICHMENTS FOR CHICKENS THAT MEET THE REQUIREMENTS OF SECTION 4.8 – SECTION A *continued*

1 enrichment = at least one item from the enrichments listed in this section (See also Section 4.8 for further details on number of enrichments by size of house)

Edible, destructible hangers



Photo: hanging eucalyptus branches



Photo: <http://featherwel.org/feedenrichments/peckingobjects>

What is the product:

Hanging plants or small bales.

Why is it an enrichment:

These materials are edible and are something that the chickens can peck at and manipulate.

What are the criteria for acceptable use:

The hanging bales or plants need to be hung at head height and be replaced regularly.

ENRICHMENTS FOR CHICKENS THAT MEET THE REQUIREMENTS OF SECTION 4.8 – SECTION A *continued*

1 enrichment = at least one item from the enrichments listed in this section (See also Section 4.8 for further details on number of enrichments by size of house)

<p>Structures – indestructible boxes</p>		<p>What is it: Wooden or plastic boxes that birds can get in or perch on.</p> <p>Why is it an enrichment: Chickens will utilize these types of structures mostly to escape from more dominant or aggressive chickens (i.e. as ‘blinds’), to sleep/rest against, and to a limited extent, for exercise.</p> <p>What are the criteria for acceptable use: Any indestructible box has to be similar in size to a straw bale (minimum of 2.5 ft × 1.5 ft) so that multiple chickens can use it at one time, when at market weight.</p>
<p>Structures - destructible boxes</p>		<p>What is it: Cardboard boxes.</p> <p>Why is it an enrichment: Chickens will utilize these types of structures mostly to escape from more dominant or aggressive chickens (i.e. as ‘blinds’), to sleep/rest against or underneath, and to a limited extent, for exercise.</p> <p>What are the criteria for acceptable use: Any destructible box has to be similar in size to a straw bale (minimum of 2.5 ft × 1.5 ft) so that multiple chickens can use it at one time, and must be strong enough to support birds that choose to sit on top of it.</p> <p>The box must be tall enough, and must have a way for birds to easily get underneath it (e.g. the hole cut in the box at the front in the picture to the left) so that chickens can go inside until at least one week prior to slaughter.</p> <p>The box must be replaced if it collapses or is otherwise destroyed by the birds.</p>

ENRICHMENTS FOR CHICKENS THAT MEET THE REQUIREMENTS OF SECTION 4.8 – SECTION A *continued*

1 enrichment = at least one item from the enrichments listed in this section (See also Section 4.8 for further details on number of enrichments by size of house)

Structures - ramps



What is it:

Wooden, plastic or plastic-coated metal ramps.

Why is it an enrichment:

Chickens will utilize these types of structures mostly to escape from more dominant or aggressive chickens (i.e. as 'blinds'), to sleep/rest against or underneath, and to a limited extent, for exercise.

What are the criteria for acceptable use:

Any ramp has to be similar in size to a straw bale (minimum of 2.5 ft × 1.5 ft) so that multiple chickens can use it at one time, when at market weight.

Ramps must have slatted or mesh tops AND the slope of the ramp must be shallow enough so that chickens can easily walk up and perch on them until they are at least one week prior to slaughter.

Ramps must be stable and allow multiple birds at market weight to use them.

The height of a ramp must allow multiple birds to sit underneath them until they are at least one week prior to slaughter.

ENRICHMENTS FOR CHICKENS THAT MEET THE REQUIREMENTS OF SECTION 4.8 – SECTION A *continued*

1 enrichment = at least one item from the enrichments listed in this section (See also Section 4.8 for further details on number of enrichments by size of house)

**Structures –
ramps**
Continued



See comments above for criteria for acceptable use

**Structures -
half barrels
and tunnels**
(turkeys
pictured in
top photo)



See comments below for criteria for acceptable use

ENRICHMENTS FOR CHICKENS THAT MEET THE REQUIREMENTS OF SECTION 4.8 – SECTION A *continued*

1 enrichment = at least one item from the enrichments listed in this section (See also Section 4.8 for further details on number of enrichments by size of house)

Structures - half barrels and tunnels *Continued*



What is it:

Plastic barrels or large pipes laid horizontally so chickens can enter the tube or sit on top of it. Or tunnels constructed of other materials.

Why is it an enrichment:

Chickens will utilize these types of structures mostly to escape from more dominant or aggressive chickens (i.e. as 'blinds'), to sleep/rest against/inside, and to a limited extent, for exercise.

What are the criteria for acceptable use:

Any barrel or pipe has to be a minimum of 2.5 ft. long and tall enough that birds can get inside until at least one week prior to slaughter, but not so tall that birds are discouraged from sitting on top of it

A pipe or barrel that is a circle in cross section (rather than cut in half to give a semi-circular cross section) must be secured in the litter (e.g. partly buried) so it does not roll when chickens go inside.

Any pipe or barrel that is a circle in cross section must either have a corrugated or similar surface that birds can grip (see picture, left), or must have litter inside the pipe.

See table on enrichments for chickens that do not meet the requirements of Section 4.8 for uses of barrels that would not count as an enrichment.

ENRICHMENTS FOR CHICKENS THAT MEET THE REQUIREMENTS OF SECTION 4.8 – SECTION B

1 enrichment = at least two (2) items (either 2 of the same, or one (1) of one type and 1 of another) from the enrichments listed in this section
(See also Section 4.8 for further details on number of enrichments by size of house)

Edible pecking blocks



What is it:

Hanging edible blocks for chickens to peck at.

Why is it an enrichment:

Chickens enjoy a wide range of foods and have an extremely strong drive to forage, scratch, and peck. Supplying “grain blocks” or “mineral pecking blocks” allows chickens to actively engage in seeking out food in a more natural way.

(NOTE: Any foodstuffs given to the birds, including enrichments, must meet the G.A.P. standards, which prohibit mammalian, avian (including eggs), and fish by-products. See Standards 3.4.1 to 3.4.3.

What are the criteria for acceptable use:

Blocks must be replaced as chickens eat them.

NOTE: Concrete blocks are not acceptable as pecking blocks.

ENRICHMENTS FOR CHICKENS THAT MEET THE REQUIREMENTS OF SECTION 4.8 – SECTION B *continued*

1 enrichment = at least two (2) items (either 2 of the same, or one (1) of one type and 1 of another) from the enrichments listed in this section
(See also Section 4.8 for further details on number of enrichments by size of house)



Bundles of strings/ropes and paper twists

Source: <https://www.cambridge.org/core/journals/animal/article/influence-of-providing-perches-and-string-on-activity-levels-fearfulness-and-leg-health-in-commercial-broiler-chickens/22316C47DA193FA25D4BFB7A3F7FCAB5#fndtn-supplementary-materials>



Photo: <http://www.featherwel.org/feedenrichments/peckingobjects>

What is the product:

Hanging inedible items for chickens to peck at.

Why is it an enrichment:

Chickens have a desire to investigate and peck at items in their environment. String is one of the simplest pecking objects used on farms. In addition, paper twists can encourage exploratory behavior in chickens and stimulate activity levels.



What are the criteria for acceptable use:

String or rope made of natural fibers is preferred. The individual strands of string or twists of paper must be presented in a bundle that is at least 1" (2.5cm) in diameter when it is first given to the chickens.

Paper twists must not be made from newspaper, magazines or any other paper containing dyes.

ENRICHMENTS FOR CHICKENS THAT MEET THE REQUIREMENTS OF SECTION 4.8 – SECTION B *continued*

1 enrichment = at least two (2) items (either 2 of the same, or one (1) of one type and 1 of another) from the enrichments listed in this section
(See also Section 4.8 for further details on number of enrichments by size of house)

<p>Small edible hangers</p>		<p>What is the product: Hanging edible items for chickens to peck at.</p> <p>Why is it an enrichment: Given chickens' strong desire to investigate, explore, peck and forage, edible hangers are also effective enrichments. Suspending broccoli, heads of lettuce, corncobs, cabbages, alfalfa or oat hay bunches, and other foods has the added benefit of stimulating physical activity.</p> <p>What are the criteria for acceptable use: The edible item needs to be hung at chicken head height and must be replaced as soon as they are consumed.</p>
<p>Shelters</p>		<p>What is it: An object that allows a bird to go underneath it to shelter.</p> <p>Why is it an enrichment: Chickens will utilize these types of structures mostly to escape from more dominant or aggressive chickens (i.e. as 'blinds'), to sleep/rest against or underneath.</p> <p>What are the criteria for acceptable use: Any shelter should be at least 2.5 ft long.</p> <p>The height of a shelter must allow multiple birds to sit underneath them until they are at least a week prior to slaughter.</p>

ENRICHMENTS FOR CHICKENS THAT MEET THE REQUIREMENTS OF SECTION 4.8 – SECTION B *continued*

1 enrichment = at least two (2) items (either 2 of the same, or one (1) of one type and 1 of another) from the enrichments listed in this section
(See also Section 4.8 for further details on number of enrichments by size of house)

Shelters *Continued*





See comments above for criteria for acceptable use

Note: The wooden shelter in the picture to the left may look, at first glance, like the “Structures – ramps” that are listed in Section B above.

However, the surface of the ramp is smooth – not meshed or slatted as required in Section A. Chickens might perch on this shelter when they are young, but will not continue to do so at older ages, as the smooth surface does not offer sufficient traction. This is therefore classified only as a shelter and not a ramp.



ENRICHMENTS FOR CHICKENS THAT DO NOT MEET THE REQUIREMENTS OF SECTION 4.8

Enrichments listed in this table do not count towards the requirements of Section 4.8; but operations may choose to use them in addition to enrichments selected from the tables above. For example, some operations like to play music to mask background noise that might startle the birds, and this can still be done, but it would not count as an enrichment to meet the requirements of Section 4.8.

<p>Non-edible hangers</p>		<p>What is the product: CDs, aluminum cans, plastic bottles, pieces of rubber hose or PVC tubing, plastic colored keys.</p> <p>Why isn't it suitable as an enrichment: While these non-edible hangers may hold the birds' interest initially, chickens of all ages quickly become indifferent to these types of hanging objects and materials that cannot be consumed.</p>
<p>Plastic barrels cut in half with solid ends</p>		<p>What is the product: Plastic drums (e.g. 55 gallon drums) that are cut in half, with the ends left on.</p> <p>Why doesn't it meet the requirements of Section 4.8: Although chickens might in theory get into barrels when placed cut side up; or get on top of barrels when placed cut side down, the smoothness of the curved sides will not make this attractive, and small birds could get trapped inside.</p> <p>If the ends of the drums were removed, they could possibly be used as tunnels and meet the requirements of Section 4.8 – see Section B above.</p>
<p>Music and radio</p>	<p><i>No photo available</i></p>	<p>What is the product: Leaving a radio or music playing in the background.</p> <p>Why doesn't it meet the requirements of Section 4.8: Although some believe that playing music or the radio can be interesting to chickens, neither specifically encourages chickens to perform natural behavior and is not considered to be an enrichment for the purposes of Section 4.8.</p>

ENRICHMENTS FOR CHICKENS **THAT DO NOT MEET THE REQUIREMENTS OF SECTION 4.8** *continued*

Enrichments listed in this table do not count towards the requirements of Section 4.8; but operations may choose to use them in addition to enrichments selected from the tables above. For example, some operations like to play music to mask background noise that might startle the birds, and this can still be done, but it would not count as an enrichment to meet the requirements of Section 4.8.

Feeders and waterers	<p><i>No photo available</i></p>	<p>Feeders and waterers are designed exclusively for maintaining nutrition and hydration, and do not serve as enrichments. Using feeder lids/egg flats filled with grain does not count as an acceptable enrichment. Certainly birds may hop onto feeders or waterlines, but these essentials are not considered to be enrichments that allow birds to engage in a broader range of natural behavior.</p>
Visual Barriers	<p><i>No photo available</i></p>	<p>Barriers include plywood sheets, shade cloth or similar material hung vertically. While the use of these barriers can provide places for birds to hide from other birds they do not encourage natural behaviors such as foraging and pecking and are not things that chickens can interact with in any way.</p>
Diatomaceous earth		<p>Some producers may give diatomaceous earth (also known as diatomite or D.E.), a light-colored, soft, and friable sedimentary rock, to chickens in their litter or in feed mixtures. However, this is not an enrichment for the purposes of Section 4.8 as it does not stimulate the birds' environment or activity levels.</p>
Litter and dust-baths (laying hen pictured)	 <p><i>Photo: https://www.rspca.org.au/free-range-standards</i></p>	<p>What is the product: An area of dry, friable material where chickens can dust-bathe.</p> <p>Why isn't it suitable as an enrichment: Although chickens will forage in litter and make use of dust-baths, such materials—whether sawdust, wood shavings, rice (or other) hulls, long or chopped straw, hay, miscanthus, sand, or chopped corn stalks—are requirements in the 5-Step® standards, rather than considered additions that enhance the chickens' environment.</p>

ENRICHMENTS FOR CHICKENS **THAT DO NOT MEET THE REQUIREMENTS OF SECTION 4.8** *continued*

Enrichments listed in this table do not count towards the requirements of Section 4.8; but operations may choose to use them in addition to enrichments selected from the tables above. For example, some operations like to play music to mask background noise that might startle the birds, and this can still be done, but it would not count as an enrichment to meet the requirements of Section 4.8.

Weigh scales



Photo: <http://www.opticonagri.com/sites/default/files/portfolio/bird-scales-weighingbreeder-broilers.jpg>

What is the product:

Weigh scales

Why isn't it suitable as an enrichment:

The weigh scale is only ~1" off the ground so that it can easily weigh birds as they move around the barn. While the number of times a scale weighs birds may be a good indicator of the overall activity of a flock, it is not considered an enrichment.

Part structures



What is the product:

Plastic frames or stools

Why isn't it suitable as an enrichment:

None of the items pictured encourages chickens to perform natural behavior and is not considered to be an acceptable enrichment.

ENRICHMENTS FOR CHICKENS **THAT DO NOT MEET** THE REQUIREMENTS OF SECTION 4.8 *continued*

Enrichments listed in this table do not count towards the requirements of Section 4.8; but operations may choose to use them in addition to enrichments selected from the tables above. For example, some operations like to play music to mask background noise that might startle the birds, and this can still be done, but it would not count as an enrichment to meet the requirements of Section 4.8.

Part structures continued



What is the product:

Plastic frames or stools

Why isn't it suitable as an enrichment:

None of the items pictured encourages chickens to perform natural behavior and is not considered to be an acceptable enrichment.

Appendix IX: Illustration of “At Chicken Height”


For vegetation to be acceptable as vegetative cover (see also Section 5.2) it must be available at “chicken height”. This is illustrated below.

Note: Vegetation in Picture 3 is inaccessible at chicken height and therefore not eligible to meet the requirements of Section 5.2. However this tall vegetation is acceptable as a provision to encourage chickens to go onto pasture, if it is within 15 ft (4.5 m) of the house. See Section 5.4.


Picture 1: Chicken can forage in grass or other short plants. Vegetation is accessible at chicken height.



Picture 2: Chicken can reach leaves on shrubs or tall grasses without having to jump or fly up. Vegetation is accessible at chicken height.



Picture 3: Chicken cannot reach leaves on shrubs or tall grasses without having to jump or fly up. Vegetation is NOT accessible at chicken height and could not be included as vegetative cover under Section 5.2.



Appendix X: Provisions to Encourage Chickens to Use Pasture

TREE COVER

Wooded areas provide acceptable cover.
ACCEPTABLE.



Young trees planted on pasture (arrow). The young trees DO NOT provide enough cover yet, so the operation would need to provide additional provisions. **NOT ACCEPTABLE.**



Mature trees providing enough cover to meet the standard. **ACCEPTABLE.**



Large, leafy, bushy trees provide acceptable cover.
ACCEPTABLE.



These young saplings will provide cover in a few years – but in the meantime the operation needs to provide additional provisions. **NOT ACCEPTABLE** unless more manmade structures are added.



These trees are close to the house and provide cover along the whole side of the house.
ACCEPTABLE.



ARTIFICIAL COVER

Shade cloth is an acceptable provision. **ACCEPTABLE.**



This manmade structure is an acceptable Provision as long as there are enough to meet Standard 5.4.3. **ACCEPTABLE.**



These A-frames are acceptable but there needs to be enough of them to meet Standard 5.4.3. **ACCEPTABLE.**



This arc is acceptable but there need to be enough of them to meet Standard 5.4.3. **ACCEPTABLE.**



This structure is acceptable but there need to be enough of them to meet Standard 5.4.3. **ACCEPTABLE.**



This structure is acceptable but there need to be enough of them to meet Standard 5.4.3. Note the bushes and shrubs also provide additional cover. **ACCEPTABLE.**



TALL PLANT COVER

While this pasture meets the vegetative cover requirements, these grasses are not tall enough.

NOT ACCEPTABLE.



Tall corn stalks provide overhead cover. However, corn stalks may not last year-round and additional provisions may need to be provided during the winter to meet the standard.

ACCEPTABLE.



Cover that is tall and dense, allowing birds to hide and explore the whole pasture.

ACCEPTABLE.



Tall, leafy plants provide overhead cover.

ACCEPTABLE.



Dense brambles provide cover opportunities even in winter.

ACCEPTABLE.



These rushes are not tall enough to provide cover.

NOT ACCEPTABLE.



Glossary

Term	Definition
Brooding	<i>Special provision of food, water and warmth for young chicks.</i>
By-product	<i>Animal waste and products derived from slaughter/harvest process including blood or any of it's components, meat, bone, bristles, flesh, hair, hides, hooves, horns, offal, skins, wool, fat, and/or feathers.</i> <i>Fish includes whole fish, parts of fish, fish meal, fish by-products from the processing industry and other aquatic species and/or products (does not include seaweed or oyster shell).</i>
Cull	<i>A bird that has been removed by a caretaker and killed as a management decision.</i>
Environmental enrichment	<i>Materials that are provided to chickens to add complexity to their environment, encourage the expression of natural behavior (such as ground scratching, pecking, and foraging), and decrease the expression of abnormal and deleterious behavior.</i>
Euthanasia	<i>The act of killing individual chickens on-farm in response to an irrecoverable illness or injury.</i>
Flock	<i>A barn/house of chickens. The group can be kept together or divided into smaller groups but would be considered one flock. See also Program Requirements 1e.</i>
Genetically modified	<i>Chickens who have been genetically altered (modified, engineered). This does not include chickens that are genetically selected for certain traits.</i>
Lameness	<i>The inability to use one or both legs in a normal manner.</i>
Litter	<i>Bedding materials.</i>
Loading	<i>Putting chickens into transport crates/containers whether manually or mechanically.</i>
Mobile coop	<i>A structure that is both moveable and moved, where chickens are confined within the structure at all times. Also known as chicken tractors.</i>
Mobile housing	<i>A structure that is both moveable and moved, at a minimum, between flocks of chickens. Mobile housing also allows chickens access to pasture.</i>
Mobile slaughter	<i>A slaughter facility that travels from operation to operation so chickens may be slaughtered on-farm.</i>
Off-label / extra-label medication	<i>Off-label use is the use of pharmaceutical drugs for an unapproved indication or in an unapproved age group, unapproved dosage, or unapproved form of administration.</i>
Operation	<i>A farm keeping chickens (see definition under Program requirements 1.f).</i>
Organophosphates	<i>Chemical compounds often used as pesticides, which have been shown to have adverse effects on the nervous system of humans and animals.</i>
On-farm slaughter facility	<i>A permanent structure or building located on the farm; designed to slaughter and process chickens.</i>
Pasture	<i>Access to rangeland, grassland, planted pastures, managed pastures, wooded areas, and any other land where chickens have access to vegetation.</i>
Placement	<i>The act of removing birds from the transport container and placing into their living accommodation.</i>
Roost	<i>The place where chickens congregate to rest at night.</i>
Sub-therapeutic	<i>Administering treatment in a preventative manner when chickens are not sick; this includes low doses of medication over an extended period of time or using medication routinely.</i>
Veranda	<i>A roofed structure, without solid side walls, along the outside of a house, level with the ground floor.</i>